arXiv:1205.0185v1 [math.RT] 1 May 2012

GRADED CARTAN DETERMINANTS OF THE SYMMETRIC GROUPS

SHUNSUKE TSUCHIOKA

Abstract. We give the graded Cartan determinants of the symmetric groups. Based on it, we propose a gradation of Hill's conjecture which is equivalent to Külshammer-Olsson-Robinson's conjecture on the generalized Cartan invariants of the symmetric groups.

1. Introduction

Let $\mathfrak{g}_0$ be a finite-dimensional simple simply-laced complex Lie algebra. The purpose of this paper is to give the Shapovalov determinants of the basic representation $V(\Lambda_0)$ of the untwisted quantum affinization $U_v(\widehat{\mathfrak{g}}_0)$ weight space wise (see Theorem 5.6) and apply them to the graded representation theory of the symmetric groups (see §6 and §7).

1.1. **Kashiwara's problem on the specialization for quantum groups.** The motivation of this paper comes from a problem of Kashiwara. Let $\mathfrak{g}$ be a symmetrizable Kac-Moody Lie algebra and let $\lambda \in \mathcal{P}^+$ be a dominant integral weight. In [Kas, Problem 2], Kashiwara asks at which specialization $v = \xi \in \mathbb{C}^\times$ (see Remark 4.14) the integrable highest weight $U_v(\mathfrak{g})$-module $V(\lambda)$ remains irreducible. When $\mathfrak{g}$ is a finite-dimensional simple complex Lie algebra, it is known that the specialized module remains irreducible if ξ is not a root of unity [APW, Theorem 6.4]. More precisely, for an ℓ-th root of unity $\xi = \exp(\frac{2\pi\sqrt{-1}}{\ell})$ where $\ell \geq 1$, the specialized $U_v(\mathfrak{g})|_{v=\xi}$-module $V(\lambda)|_{v=\xi}$ is irreducible if $\ell < (\alpha, \lambda)$ for any positive root $\alpha \in \Delta^+$ of $\mathfrak{g}$ [APW, Theorem 6.7].

When $\mathfrak{g}$ is an affine Kac-Moody Lie algebra, almost nothing is known about Kashiwara's problem. Chari-Jing proved that the specialized $U_v(\widehat{\mathfrak{g}}_0)|_{v=\xi}$-module $V(\Lambda_0)|_{v=\xi}$ is irreducible if ξ is not a root of unity. More precisely, for an ℓ-th root of unity $\xi = \exp(\frac{2\pi\sqrt{-1}}{\ell})$ where $\ell \geq 1$, if ℓ is coprime to the Coxeter number N of $\mathfrak{g}_0$, then the specialized module is irreducible [CJ, Theorem 4]. Recall that $N = n$ (resp. $2(m-1), 12, 18, 30$) when $\mathfrak{g}_0$ is of type A_{n-1} for $n \geq 2$ (resp. D_m for $m \geq 4$, E_6, E_7, E_8). The main result of this paper (Theorem 5.6) gives an answer to Kashiwara's problem when $\mathfrak{g} = \widehat{\mathfrak{g}}_0$ and λ is level 1.

Theorem 1.1. *Let $\mathfrak{g}_0$ be a finite-dimensional simple simply-laced complex Lie algebra of type X. Then, for $\ell \geq 1$ the specialized $U_v(\widehat{\mathfrak{g}}_0)|_{v=\exp(\frac{2\pi\sqrt{-1}}{\ell})}$-module $V(\Lambda_0)|_{v=\exp(\frac{2\pi\sqrt{-1}}{\ell})}$ is irreducible if and only if*

(a) $\mathsf{gcd}(\ell, 2n) = 1, 2$ *when $X = A_{n-1}$ for $n \geq 2$,*

(b) $\ell \notin 4\mathbb{Z}$ *(resp. $\notin 3\mathbb{Z}, \notin 4\mathbb{Z}, \notin 60\mathbb{Z}$) when $X = D_n$ for $n \geq 4$ (resp. E_6, E_7, E_8).*

Date: May 1, 2012.

2000 *Mathematics Subject Classification.* Primary 81R50, Secondary 20C08.

Key words and phrases. symmetric groups, Iwahori-Hecke algebras, graded representation theory, quantum groups, Shapovalov forms, quiver Hecke algebras, generalized blocks.

The research was supported by Grant-in-Aid for Research Activity Startup 22840026 and Research Fellowships for Young Scientists 23·8363, Japan Society for the Promotion of Science.

1.2. **Graded Cartan determinants.** Recently, Khovanov-Lauda and Rouquier independently introduced a remarkable family of algebras (the KLR algebras) that categorifies the negative half of the quantized enveloping algebras associated with symmetrizable Kac-Moody Lie algebras [KL1, KL2, Rou] (see Definition 6.7). An application of the KLR algebras is the homogeneous presentation of the symmetric group algebras [BK2, Rou] (see Theorem 6.15) which quantize [BK1] Ariki's categorification of the Kostant $\mathbb{Z}$-form of the basic $\widehat{\mathfrak{sl}}_p$-module $V(\Lambda_0)^{\mathbb{Z}} \cong \bigoplus_{n\geq 0} \mathsf{K}_0(\mathsf{Proj}(\mathbb{F}_p\mathfrak{S}_n))$. Since the Shapovalov-like form (see Proposition 4.8) on the left hand side and the graded Cartan pairing on the right hand side are compatible [BK3] (see Theorem 6.16), the main result of this paper (Theorem 5.6) can be interpreted as the graded Cartan determinants of the symmetric groups. The story is also valid for its q-analog, the Iwahori-Hecke algebra of type A (see §6.4).

Theorem 1.2. *Let $p \geq 2$ be a prime number (resp. $p \geq 2$ an arbitrary integer). Then, the graded Cartan determinant of a block of the symmetric group algebra over characteristic p (resp. the Iwahori-Hecke algebra of type A over quantum characteristic p) whose p-weight is d is given by $\prod_{s=1}^{d}[p]_s^{N_{p,d,s}}$ where*

$$N_{p,d,s} = \sum_{\lambda\in\mathsf{Par}(d)} \frac{m_s(\lambda)}{p-1}\prod_{u\geq 1}\binom{m_u(\lambda)+p-2}{m_u(\lambda)} = \sum_{(\lambda_i)_{i=1}^{p-1}\in\mathsf{Par}_{p-1}(d)} m_s(\lambda_1).$$

For the notations on partitions, see Notations in the end of §1.

In virtue of the classical results of the modular representation theory of finite groups, the Cartan determinant (see Definition 6.1) of (a block of) a group algebra of a finite group G over characteristic $p \geq 2$ is a power of p [Bra, Theorem 1] and its degree reflects p-defects of p-regular conjugacy classes of G [BrNe, Part III,§16] (see Theorem 7.6). Based on Olsson's calculation of p-defects of the symmetric groups [Ols], Bessenrodt-Olsson gave an explicit formula for the Cartan determinants of (a block of) a symmetric group algebra over characteristic p in terms of generating functions [BO1, Theorem 3.3, Theorem 3.4].

On the other hand, Brundan-Kleshchev arrived at the same formula by a different method [BK4, Corollary 1]. They utilized Ariki's categorification and turned the calculation of the p-Cartan determinants of the symmetric groups into the calculation of the Shapovalov determinants of the basic $\widehat{\mathfrak{sl}}_p$-module $V(\Lambda_0)$. Their block-theory-free method has an advantage in that it is applicable to the Iwahori-Hecke algebra of type A. In fact, Brundan-Kleshchev settled affirmatively the conjecture of Mathas that the Cartan determinants of (a block of) the Iwahori-Hecke algebra of type A over quantum characteristic ℓ is a power of ℓ for arbitrary $\ell \geq 2$ (see [Don]).

Theorem 1.2 (and Theorem 5.6) can be seen a gradation of Brundan-Kleshchev's result and our method is similar to theirs. A difference is that we use a bi-additive form that obeys a rule slightly different from that of the Shapovalov form (see Proposition 4.8).

We remark that Brundan-Kleshchev also calculated the Shapovalov determinants of the basic module over a twisted affinization of $\mathfrak{g}_0$. Although our method should be applicable in the quantum setting, there are some obstructions to overcome (see Remark 5.11). We discuss them and give a conjectural Shapovalov determinants (see Conjecture 5.13).

Recently, the authors [KKT] introduced a superversion of the KLR algebras which they call quiver Hecke superalgebras and showed "Morita superequivalence" between the spin symmetric group algebras and quiver Hecke superalgebras [KKT, Theorem 5.4]. From the results of [BK5, Tsu], it is expected that the quiver Hecke superalgebras categorify the quantum groups (see also [HW]) and through this the Shapovalov determinants of the basic module of $U_v(A^{(2)}_{p-1})$ for odd prime $p \geq 3$ coincides with the graded Cartan determinants of the spin symmetric group algebras.

1.3. Külshammer-Olsson-Robinson's conjecture and graded Cartan invariants.

Based on an observation that combinatorial notions for partitions (such as p-cores) that appear in the description of the p-modular representation-theoretic invariants of the symmetric groups make sense when p is not a prime, Külshammer-Olsson-Robinson initiated the study of "ℓ-modular representations of the symmetric groups" for arbitrary $\ell \geq 2$ [KOR].

Their main result is a generalization of Nakayama-Brauer-Robinson classification of the p-blocks of $\mathfrak{S}_n$ for an any $\ell \geq 2$ [KOR, Theorem 5.13] (see Theorem 7.4). Based on it, they gave a conjectural generalization of the Cartan invariants of the symmetric groups for an arbitrary $\ell \geq 2$ [KOR, Conjecture 6.4] (see Conjecture 7.10 which we call the KOR conjecture for short). As an evidence, they checked that their invariants gives a generalized Cartan determinant calculated by Brundan-Kleshchev [BK4].

Hill settled affirmatively the KOR conjecture for any $\ell \geq 2$ such that each prime factor p of ℓ divides ℓ at most p times [Hil, Theorem 1.3]. In his course of proof, Hill gave a conjecture [Hil, Conjecture 10.5] (see Conjecture 7.14). Hill's conjecture and the KOR conjecture are equivalent in virtue of [BH] (see Corollary 7.16).

As graded Cartan determinants calculated in this paper, it is reasonable to expect that a proof of Hill's conjecture similarly works in a graded setting. Motivated by this, we propose a gradation of Hill's conjecture (see Conjecture 7.17). As a support, we check that Conjecture 7.17 is compatible with the graded Cartan determinants (see Theorem 7.20). The proof itself may be of interest. We expect that our gradation is correct and gives an insight to future trials of the proof of Hill's conjecture (and thus the KOR conjecture).

Notations. $\mathbb{N} = \mathbb{Z}_{\geq 0}$ (resp. $\mathbb{N}_+ = \mathbb{Z}_{\geq 1}$) means the set of non-negative (resp. positive) integers. We denote the empty partition by ϕ and reserve the symbol $\emptyset$ for the empty set. For a partition $\lambda = (\lambda_1, \lambda_2, \cdots)$, we define $m_k(\lambda) = |\{i \geq 1 \mid \lambda_i = k\}|$ for $k \geq 1$. We also define $|\lambda| = \sum_{i\geq 1} \lambda_i$ and $\ell(\lambda) = \sum_{i\geq 1} m_i(\lambda)$. We denote by $\mathsf{Par}(n)$ the set of partitions of $n \geq 0$ and define $\mathsf{Par} = \bigsqcup_{n\geq 0} \mathsf{Par}(n)$. For $m, n \geq 0$, we denote by $\mathsf{Par}_m(n)$ the multipartitions of n, i.e., $\mathsf{Par}_m(n) = \{(\lambda_i)_{i=1}^m \in \mathsf{Par}^m \mid \sum_{i=1}^m |\lambda_i| = n\}$ and put $u(m, n) = |\mathsf{Par}_m(n)|$. Note that $u(0, 0) = 1$ and $u(0, n) = 0$ for $n \geq 1$.

Organization of the paper In §2, we develop a linear algebra for Gram determinants. In §3, we review the boson-fermion correspondence and deduce the existence of a bi-additive form we need. In §4, we give a brief review on quantum groups. In §5, we calculate the Shapovalov determinants of the basic representations of the untwisted affine A,D,E quantum groups and in §6 using the KLR algebras we interpret it as graded Cartan determinants of the symmetric groups and its Iwahori-Hecke akgebras. Finally, in §7 we propose a gradation on Hill's conjecture and make some justifications.

Acknowledgements. The content of §7 is benefited from the discussion with Yasuhide Numata. The author would like to thank him and also would like to thank Hiraku Nakajima and Yoshiyuki Kimura for discussions on quantum groups. The author is grateful to Hiro-Fumi Yamada. Without his encouragement and their paper [ASY], this paper would not have been written.

2. Linear Algebra on Gram determinants

Let $\mathbf{k}$ be a field and let V be a $\mathbf{k}$-vector space. We say that a map $B : V \times V \to \mathbf{k}$ is bi-additive if we have $B(w_1 + w_2, w) = B(w_1, w) + B(w_2, w)$ and $B(w, w_1 + w_2) = B(w, w_1) + B(w, w_2)$ for all $w, w_1, w_2 \in V$. The radical $\mathsf{Rad}(B)$ of a bi-additive map B is defined to be one of the following two additive subgroups of V

$$\{w_1 \in V \mid \forall w_2 \in V, B(w_1, w_2) = 0\}, \quad \{w_2 \in V \mid \forall w_1 \in V, B(w_1, w_2) = 0\}$$

if they are equal (otherwise, we do not define $\mathsf{Rad}(B)$). We say that a bi-additive map B is non-degenerate if $\mathsf{Rad}(B)$ is defined and $\mathsf{Rad}(B) = 0$.

Definition 2.1. *Let* $(\mathbf{k}, \tau, \mathscr{A}, V, S, V^{\mathscr{A}})$ *be a 6-tuple such that*

(a) $\mathbf{k}$ *is a field with a ring involution* $\tau : \mathbf{k} \xrightarrow{\sim} \mathbf{k}$,
(b) $\mathscr{A}$ *is a subring of* $\mathbf{k}$ *such that* $\tau(\mathscr{A}) \subseteq \mathscr{A}$,
(c) V *is a finite dimensional* $\mathbf{k}$*-vector space,*
(d) $V^{\mathscr{A}}$ *is an* $\mathscr{A}$*-lattice of* V*, i.e.,* $V^{\mathscr{A}}$ *is a free* $\mathscr{A}$*-module with an isomorphism* $\mathbf{k} \otimes_{\mathscr{A}} V^{\mathscr{A}} \xrightarrow{\sim} V$,
(e) $S : V \times V \to \mathbf{k}$ *is a bi-additive non-degenerate map such that* $S(aw_1, w_2) = \tau(a)S(w_1, w_2)$ *and* $S(w_1, aw_2) = aS(w_1, w_2)$ *for all* $w_1, w_2 \in V$ *and* $a \in \mathbf{k}$.

To such a 6-tuple, we define the Gram matrix $\mathsf{GM}_{\mathbf{k},\tau,\mathscr{A}}(V, S, V^{\mathscr{A}}) = (S(w_i, w_j))_{1 \leq i,j \leq \dim V}$ *and define the Gram determinant* $\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V, S, V^{\mathscr{A}}) = \det \mathsf{GM}_{\mathbf{k},\tau,\mathscr{A}}(V, S, V^{\mathscr{A}})$ *where* $(w_i)_{1 \leq i \leq \dim V}$ *is an* $\mathscr{A}$*-basis of* $V^{\mathscr{A}}$.

Let X and Y are two Gram matrices associated with two choices of $\mathscr{A}$-bases of $V^{\mathscr{A}}$. Clearly, there exists $P \in \mathrm{GL}_{\dim V}(\mathscr{A})$ such that $X = \tau({}^{\mathrm{tr}}P)YP$. Thus, $\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V, S, V^{\mathscr{A}})$ is uniquely determined as an element of $\mathbf{k}^{\times}/\{a\tau(a) \mid a \in \mathscr{A}^{\times}\}$.

The following comparison result is based on [BK4, §5].

Proposition 2.2. *Let* I *be a finite set and let* $\mathbf{k}$ *be a field. We regard a polynomial ring* $V = \mathbf{k}[y^{(i)}_n \mid i \in I, n \geq 1]$ *as a graded* $\mathbf{k}$*-algebra via* $\mathsf{deg}\, y^{(i)}_n = n$ *and denote by* V_d *the* $\mathbf{k}$*-vector subspace of* V *consisting of homogeneous elements of degree* d *for* $d \geq 0$*. Assume* $\operatorname{char} \mathbf{k} = 0$ *and we are given the following data.*

(a) *a subring* $\mathscr{A}$ *of* $\mathbf{k}$ *and a ring involution* $\tau : \mathbf{k} \xrightarrow{\sim} \mathbf{k}$ *such that* $\tau(\mathscr{A}) \subseteq \mathscr{A}$,
(b) *a family of invertible matrices* $A = (A^{(m)})_{m \geq 1}$ *where* $A^{(m)} = (a^{(m)}_{ij})_{i,j \in I} \in \mathrm{GL}_I(\mathscr{A}^{\tau})$,
(c) *() two bi-additive non-degenerate maps* $\langle , \rangle_S, \langle , \rangle_K : V \times V \to \mathbf{k}$ *such that*
    - $\langle af, g \rangle_X = \tau(a)\langle f, g \rangle_X$, $\langle f, ag \rangle_X = a\langle f, g \rangle_X$ *and* $\langle f, g \rangle_X = \tau(\langle g, f \rangle_X)$
    - $\langle 1, 1 \rangle_S = \langle 1, 1 \rangle_K$ *and* $\langle m y^{(i)}_m f, g \rangle_S = \langle f, \sum_{j \in I} a^{(m)}_{ij} \frac{\partial g}{\partial y^{(j)}_m} \rangle_S$, $\langle m y^{(i)}_m f, g \rangle_K = \langle f, \frac{\partial g}{\partial y^{(i)}_m} \rangle_K$,

    for $X \in \{S, K\}$ *and* $f, g \in V, a \in \mathbf{k}, m \geq 1$,
(d) *a family of new variables* $(x^{(i)}_n)_{\substack{i \in I,\\ n \geq 1}}$ *such that* $x^{(i)}_n - y^{(i)}_n \in \mathbf{k}[y^{(i)}_m \mid 1 \leq m < n] \cap V_n$.

Then, in $\mathbf{k}^{\times}/\{a\tau(a) \mid a \in \mathscr{A}^{\times}\}$ *we have*

$$\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V_d, \langle , \rangle_S, V^{\mathscr{A}}_d)/\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V_d, \langle , \rangle_K, V^{\mathscr{A}}_d) = \Delta_d(A)$$

for any $d \geq 0$ *where* $V^{\mathscr{A}}_d = V_d \cap \mathscr{A}[x^{(i)}_n \mid i \in I, n \geq 1]$ *and*

$$\Delta_d(A) = \prod_{s=1}^{d} (\det A^{(s)})^{\sum_{\lambda \in \mathsf{Par}(d)} \frac{m_s(\lambda)}{|I|} \prod_{u \geq 1} \binom{m_u(\lambda)+|I|-1}{m_u(\lambda)}}.$$

Proof. Firstly, we note that in virtue of (d) we have $V = \mathbf{k}[x_n^{(i)} \mid i \in I, n \geq 1]$. Thus, $V_d^{\mathscr{A}}$ is an $\mathscr{A}$-lattice of V_d.

From now on, we fix a total ordering on I and define

$$\Omega(\lambda) = \{(i_1, \cdots, i_{\ell(\lambda)}) \in I^{\ell(\lambda)} \mid \lambda_j = \lambda_{j+1} \Rightarrow i_j \geq i_{j+1}\}$$

for $\lambda \in \mathsf{Par}$ and put $\Omega_d = \{(\lambda, i) \mid \lambda \in \mathsf{Par}(d), i \in \Omega(\lambda)\}$ for $d \geq 0$.

We define a family of new variables $z_n^{(i)} = \sum_{j \in I} a_{ij}^{(n)} y_n^{(j)}$ for $i \in I$ and $n \geq 1$. Then, we see that for any $d \geq 0$

- $\{x_\lambda^{(i)} := \prod_{k=1}^{\ell(\lambda)} x_{\lambda_k}^{(i_k)}\}_{(\lambda,i)\in\Omega_d}$ is an $\mathscr{A}$-basis of $V_d^{\mathscr{A}}$,
- $\{y_\lambda^{(i)} := \prod_{k=1}^{\ell(\lambda)} y_{\lambda_k}^{(i_k)}\}_{(\lambda,i)\in\Omega_d}$ and $\{z_\lambda^{(i)} := \prod_{k=1}^{\ell(\lambda)} z_{\lambda_k}^{(i_k)}\}_{(\lambda,i)\in\Omega_d}$ are $\mathbf{k}$-basis of V_d.

We show for any $\lambda \in \mathsf{Par}$ by induction on $\ell := \ell(\lambda) \geq 0$ that we have

$$\forall i \in \Omega(\lambda), \forall f \in V, \langle y_\lambda^{(i)}, f\rangle_S = \langle z_\lambda^{(i)}, f\rangle_K. \tag{2.1}$$

It is clear by (c) when $\ell = 0$ and assume that (2.1) holds for any λ with $\ell(\lambda) < \ell$. Put $j = (i_2, \cdots, i_\ell)$ and $\mu = (\lambda_2, \cdots, \lambda_\ell)$ and (2.1) is proved as follows.

$$\begin{aligned}\langle y_\lambda^{(i)}, f\rangle_S = \langle y_{\lambda_1}^{(i_1)} y_\mu^{(j)}, f\rangle_S &= \frac{1}{\lambda_1}\langle y_\mu^{(j)}, \sum_{k\in I} a_{i_1,k}^{(\lambda_1)} \frac{\partial f}{\partial y_{\lambda_1}^{(k)}}\rangle_S \\ &= \frac{1}{\lambda_1}\langle z_\mu^{(j)}, \sum_{k\in I} a_{i_1,k}^{(\lambda_1)} \frac{\partial f}{\partial y_{\lambda_1}^{(k)}}\rangle_K = \langle z_{\lambda_1}^{(i_1)} z_\mu^{(j)}, f\rangle_K = \langle z_\lambda^{(i)}, f\rangle_K.\end{aligned}$$

We note that $\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V_d, \langle,\rangle_S, V_d^{\mathscr{A}}) = \det M_S$ and $\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V_d, \langle,\rangle_K, V_d^{\mathscr{A}}) = \det M_K$ where $M_S = (\langle x_\lambda^{(i)}, x_\mu^{(j)}\rangle_S)_{(\lambda,i),(\mu,j)\in\Omega_d}$ and $M_K = (\langle x_\lambda^{(i)}, x_\mu^{(j)}\rangle_K)_{(\lambda,i),(\mu,j)\in\Omega_d}$.

Let $P \in \mathrm{GL}_{\Omega_d}(\mathbf{k})$ and $Q \in \mathrm{GL}_{\Omega_d}(\mathscr{A}^\tau)$ by

$$x_\lambda^{(i)} = \sum_{(\mu,j)\in\Omega_d} p_{\lambda,\mu}^{(i,j)} y_\mu^{(j)}, \quad z_\lambda^{(i)} = \sum_{(\mu,j)\in\Omega_d} q_{\lambda,\mu}^{(i,j)} y_\mu^{(j)}.$$

In virtue of (2.1), we have

$$\begin{aligned} M_S &= \tau(P)(\langle y_\lambda^{(i)}, x_\mu^{(j)}\rangle_S)_{(\lambda,i),(\mu,j)\in\Omega_d} = \tau(P)(\langle z_\lambda^{(i)}, x_\mu^{(j)}\rangle_K)_{(\lambda,i),(\mu,j)\in\Omega_d}, \\ M_K &= \tau(P)(\langle y_\lambda^{(i)}, x_\mu^{(j)}\rangle_K)_{(\lambda,i),(\mu,j)\in\Omega_d} = \tau(P)Q^{-1}(\langle z_\lambda^{(i)}, x_\mu^{(j)}\rangle_K)_{(\lambda,i),(\mu,j)\in\Omega_d}. \end{aligned}$$

Thus, we have $M_S = \tau(P)Q\tau(P)^{-1}M_K$ and it is enough to show that $\det Q = \Delta_d$.

It is easily seen that $q_{\lambda,\mu}^{(i,j)} = 0$ for $(\lambda, i), (\mu, j) \in \Omega_d$ with $\lambda \neq \mu$. Since $(q_{\lambda,\lambda}^{(i,j)})_{i,j\in\Omega(\lambda)}$ is identified with $\otimes_{t\geq 1} \mathsf{Sym}^{m_t(\lambda)}(A^{(t)})$, we have

$$\det Q = \prod_{\lambda\in\mathsf{Par}(d)} \det((q_{\lambda,\lambda}^{(i,j)})_{i,j\in\Omega(\lambda)}) = \prod_{\lambda\in\mathsf{Par}(d)} \det(\otimes_{t\geq 1}\mathsf{Sym}^{m_t(\lambda)}(A^{(t)})).$$

Apply Lemma 2.3 which is an easy linear algebra exercise, we get

$$\begin{aligned}\det Q &= \prod_{\lambda\in\mathsf{Par}(d)} \prod_{t\geq 1} \det(\mathsf{Sym}^{m_t(\lambda)}(A^{(t)}))^{\prod_{u\neq t}\dim \mathsf{Sym}^{m_u(\lambda)}(k^I)} \\ &= \prod_{\lambda\in\mathsf{Par}(d)} \prod_{t\geq 1} (\det(A^{(t)}))^{\binom{|I|+m_t(\lambda)-1}{m_t(\lambda)-1}\prod_{u\neq t}\binom{|I|+m_u(\lambda)-1}{m_u(\lambda)}}.\end{aligned}$$

Since $\binom{|I|+m_t(\lambda)-1}{m_t(\lambda)-1}\prod_{u\neq t}\binom{|I|+m_u(\lambda)-1}{m_u(\lambda)} = \frac{m_t(\lambda)}{|I|}\prod_u \binom{|I|+m_u(\lambda)-1}{m_u(\lambda)}$, we have $\det Q = \Delta_d$. □

Lemma 2.3. *Let* $\mathbf{k}$ *be a field.*

(a) *Let* V *be an* n*-dimensional* $\mathbf{k}$*-vector space and let* $f \in \mathsf{End}_{\mathbf{k}}(V)$. *For* $m \geq 0$, *we have*

$$\dim \mathsf{Sym}^m V = \binom{n+m-1}{m}, \quad \det \mathsf{Sym}^m f = (\det f)^{\binom{n+m-1}{m-1}}.$$

(b) *For an* ℓ*-tuple of matrices* $(B^{(i)})_{1 \leq i \leq \ell}$ *with* $B^{(i)} \in \mathsf{Mat}_{n_i}(\mathbf{k})$, *we have* $\det(\otimes_{i=1}^{\ell} B^{(i)}) = \prod_{i=1}^{\ell} (\det B^{(i)})^{N/n_i}$ *where* $N = \prod_{i=1}^{\ell} n_i$.

3. Boson-fermion correspondence

Definition 3.1. *Let* R *be a commutative ring with* 1.

(a) *The fermionic Fock space* $\mathcal{F}_R = R^{\oplus \mathsf{SIM}}$ *over* R *is a free* R*-module with a basis consisting of all semi-infinite monomials*

$$\mathsf{SIM} := \{ i_1 \wedge i_2 \wedge \cdots \mid \forall k \geq 1, i_k > i_{k+1} \text{ and } \exists N > 0, \forall k > N, i_k = i_{k-1} - 1 \}.$$

(b) *The bosonic Fock space* $\mathcal{B}_R$ *over* R *is a polynomial ring* $R[v, v^{-1}, y_1, y_2, \cdots]$.

For $m \in \mathbb{Z}$, we denote by $|m\rangle$ the vacuum vector of charge m, i.e., $|m\rangle = m \wedge (m-1) \wedge \cdots \in \mathsf{SIM}$. We say that a semi-infinite monomial $i_1 \wedge i_2 \wedge \cdots \in \mathsf{SIM}$ is of charge m if it differs from $|m\rangle$ only at a finite number of places, i.e., there exists $N > 0$ such that we have $i_k = m + 1 - k$ for all $k > N$.

Recall that for $j \in \mathbb{Z}$ a creation operator $\psi_j \in \mathsf{End}_R(\mathcal{F}_R)$ and an annihilation operator $\psi_j^* \in \mathsf{End}_R(\mathcal{F}_R)$ are defined as follows where we regard i_0 as $+\infty$.

$$\psi_j(i_1 \wedge i_2 \wedge \cdots) = \begin{cases} 0 & \exists s \geq 1, j = i_s, \\ (-1)^s i_1 \wedge \cdots \wedge i_s \wedge j \wedge i_{s+1} \wedge \cdots & \exists s \geq 0, i_s > j > i_{s+1}, \end{cases}$$

$$\psi_j^*(i_1 \wedge i_2 \wedge \cdots) = \begin{cases} 0 & \forall s \geq 1, j \neq i_s, \\ (-1)^{s+1} i_1 \wedge \cdots \wedge i_{s-1} \wedge i_{s+1} \wedge \cdots & \exists s \geq 1, i_s = j. \end{cases}$$

Remark 3.2. For any $s, t \in \mathsf{SIM}$ and $j \in \mathbb{Z}$, we have $\psi_j(s) = t$ if and only if $\psi_j^*(t) = s$.

Remark 3.3. $\mathcal{B}_R$ has a natural structure of module over the Heisenberg algebra $\mathscr{H}_R := \langle \{\alpha_m\}_{m \in \mathbb{Z}} \mid \alpha_m \alpha_n - \alpha_n \alpha_m = m \delta_{m+n,0} \rangle_{R\text{-alg}}$ through the assignment of linear operators

$$\alpha_0 = v \frac{\partial}{\partial v}, \quad \alpha_k = \frac{\partial}{\partial y_k}, \quad \alpha_{-k} = k y_k$$

where $k \geq 1$. On the other hand, bosonization turns $\mathcal{F}_R$ into an $\mathscr{H}_R$-module where $\ell \neq 0$

$$\alpha_k = \sum_{j \in \mathbb{Z}} \psi_j \psi_{j+k}^*, \quad \alpha_0 = \sum_{j > 0} \psi_j \psi_j^* - \sum_{j \leq 0} \psi_j^* \psi_j.$$

Definition 3.4. *For a partition* $\lambda \in \mathsf{Par}$, *the Schur function* $s_\lambda \in \mathbb{Q}[y_1, y_2, \cdots]$ *is defined as* $s_\lambda = \det(x_{\lambda_i - i + j})_{1 \leq i,j \leq |\lambda|}$ *where* $x_m = \delta_{m,0}$ *for* $m \leq 0$ *and*

$$1 + \sum_{n \geq 1} x_n z^n = \exp(\sum_{n \geq 1} y_n z^n). \tag{3.1}$$

Proposition 3.5 ([Kac, §14.10]). *Let* $\mathbf{k}$ *be a field of characteristic 0.*

(i) *There exists a unique* $\mathscr{H}_{\mathbf{k}}$*-module isomorphism* $\sigma_{\mathbf{k}} : \mathcal{F}_{\mathbf{k}} \xrightarrow{\sim} \mathcal{B}_{\mathbf{k}}$ *such that* $\sigma_{\mathbf{k}}(|m\rangle) = v^m$ *for any* $m \in \mathbb{Z}$.

(ii) *For any* $m \in \mathbb{Z}$ *and* $\boldsymbol{i} = i_1 \wedge i_2 \wedge \cdots \in \mathsf{SIM}$ *of charge* m*, we have* $\sigma_{\mathbf{k}}(\boldsymbol{i}) = v^m s_{\lambda(\boldsymbol{i})}$ *where* $\lambda(\boldsymbol{i}) = (i_1 - m, i_2 - (m-1), \cdots) \in \mathsf{Par}$.

Corollary 3.6 ([DcKK, Theorem 2]). *Let* $\mathbf{k}$ *be a field of characteristic 0. Define a family of new variables* $(x_n)_{n\ge 1}$ *by (3.1) and put* $\widetilde{\mathcal{B}}_{\mathbb{Z}} = \mathbb{Z}[v, v^{-1}, x_1, x_2, \cdots](\subseteq \mathcal{B}_{\mathbf{k}})$*. Then, the restriction* $\sigma_{\mathbb{Z}} = \sigma_{\mathbf{k}}|_{\mathcal{F}_{\mathbb{Z}}}$ *induces an* $\mathscr{H}_{\mathbb{Z}}$*-module isomorphism* $\sigma_{\mathbb{Z}} : \mathcal{F}_{\mathbb{Z}} \xrightarrow{\sim} \widetilde{\mathcal{B}}_{\mathbb{Z}}$.

Corollary 3.7. *Let* $\mathbf{k}$ *be a field. We regard a polynomial ring* $V = \mathbf{k}[y_1, y_2, \cdots]$ *as a graded* $\mathbf{k}$*-algebra via* $\mathsf{deg}\, y_n = n$ *and denote by* V_d *the* $\mathbf{k}$*-vector subspace of* V *consisting of homogeneous elements of degree* d *for* $d \ge 0$*. Assume* $\operatorname{char} \mathbf{k} = 0$ *and take a ring involution* $\tau : \mathbf{k} \xrightarrow{\sim} \mathbf{k}$.

(a) *There exists a unique bi-additive non-degenerate map* $\langle,\rangle_K : V \times V \to \mathbf{k}$ *such that*
  (i) $\langle af, g\rangle_K = \tau(a)\langle f, g\rangle_K, \langle f, ag\rangle_K = a\langle f, g\rangle_K$ *and* $\langle f, g\rangle_K = \tau(\langle g, f\rangle_K)$,
  (ii) $\langle 1, 1\rangle_K = 1$ *and* $\langle m y_m^{(i)} f, g\rangle_K = \langle f, \frac{\partial g}{\partial y_m^{(i)}}\rangle_K$,

for any $f, g \in V$ *and* $a \in \mathbf{k}$.

(b) *Let* $\mathscr{A}$ *be a subring of* $\mathbf{k}$ *such that* $\tau(\mathscr{A}) \subseteq \mathscr{A}$*. We choose an* $\mathscr{A}$*-lattice* $V_d^{\mathscr{A}}$ *of* V_d *by* $V_d^{\mathscr{A}} = V_d \cap \mathscr{A}[x_n \mid n \ge 1]$ *for* $d \ge 0$ *where* $(x_n)_{n\ge1}$ *is defined in terms of* $(y_n)_{n\ge1}$ *via (3.1). Then, for all* $d \ge 0$ *we have* $\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V_d, \langle,\rangle_K, V_d^{\mathscr{A}}) = 1$ *in* $\mathbf{k}^\times/\{a\tau(a) \mid a \in \mathscr{A}^\times\}$.

Proof. (a) Uniqueness is clear and we prove the existence. We introduce a bi-additive non-degenerate map $(,) : \mathcal{F}_{\mathbf{k}} \times \mathcal{F}_{\mathbf{k}} \to \mathbf{k}$ as $(\sum_{s\in\mathsf{SIM}} a_s \cdot s, \sum_{t\in\mathsf{SIM}} b_t \cdot t) = \sum_{u\in\mathsf{SIM}} \tau(a_u) b_u$. Let $\mathcal{F}_{\mathbf{k}}^{(0)}$ be the $\mathbf{k}$-subspace of $\mathcal{F}_{\mathbf{k}}$ spanned by $\{s \in \mathsf{SIM} \mid s \text{ is of charge } 0\}$. Then, $\sigma_{\mathbf{k}}|_{\mathcal{F}_{\mathbf{k}}^{(0)}} : \mathcal{F}_{\mathbf{k}}^{(0)} \xrightarrow{\sim} V(\subseteq \mathcal{B}_{\mathbf{k}})$ carries $(,)|_{\mathcal{F}_{\mathbf{k}}^{(0)} \times \mathcal{F}_{\mathbf{k}}^{(0)}}$ on $\mathcal{F}_{\mathbf{k}}^{(0)}$ into the desired map $\langle,\rangle_K$ on V because of Remark 3.2 and Remark 3.3.

(b) It is well-known that $\{s_\lambda \mid \lambda \in \mathsf{Par}\}$ forms a $\mathbb{Z}$-basis of $V^{\mathbb{Z}} := \mathbb{Z}[x_1, x_2, \cdots] \subseteq V$. From the construction of $\langle,\rangle_K$ above, it is clear that $\{s_\lambda \mid \lambda \in \mathsf{Par}\}$ is orthonormal on $V^{\mathbb{Z}}$ (and thus $\mathscr{A} \otimes V^{\mathbb{Z}}$). ☐

Corollary 3.8. *In the setting of Proposition 2.2, we may take* $x_n^{(i)}$ *for* $i \in I$ *and* $n \ge 1$ *by*

$$1 + \sum_{n\ge1} x_n^{(i)} z^n = \exp(\sum_{n\ge1} y_n^{(i)} z^n).$$

Then, we have $\mathsf{Gd}_{\mathbf{k},\tau,\mathscr{A}}(V_d, \langle,\rangle_S, V_d^{\mathscr{A}}) = \Delta_d(A)$.

4. Quantum groups and Shapovalov forms

Let v be an indeterminate. In the rest of the paper, we work in the field $\mathbf{k} = \mathbb{Q}(v)$ and its subring $\mathscr{A} = \mathbb{Z}[v, v^{-1}]$. We consider two ring involution of $\mathbf{k}$. One is the identity map $\mathsf{id}_{\mathbf{k}}$ and the other is a $\mathbb{Q}$-algebra involution $\tau : \mathbf{k} \to \mathbf{k}$ which maps v to v^{-1}.

4.1. **Quantum groups.** Let $A = (a_{ij})_{i,j\in I}$ be a symmetrizable GCM and take the symmetrization $d = (d_i)_{i\in I}$ of A, i.e., a unique $d \in \mathbb{N}_+^I$ such that $d_i a_{ij} = d_j a_{ji}$ for all $i, j \in I$ and $\gcd(d_i)_{i\in I} = 1$. We take a Cartan data $(\mathcal{P}, \mathcal{P}^\vee, \Pi, \Pi^\vee)$ in the following sense.

(a) $\mathcal{P}^\vee$ is a free $\mathbb{Z}$-module of rank $(2|I| - \operatorname{rank} A)$ and $\mathcal{P} = \operatorname{Hom}_{\mathbb{Z}}(\mathcal{P}^\vee, \mathbb{Z})$,
(b) $\Pi^\vee = \{h_i \mid i \in I\}$ is a $\mathbb{Z}$-linearly independent subset of $\mathcal{P}^\vee$,
(c) $\Pi = \{\alpha_i \mid i \in I\}$ is a $\mathbb{Z}$-linearly independent subset of $\mathcal{P}$,
(d) $\alpha_j(h_i) = a_{ij}$ for all $i, j \in I$.

We denote by $\mathcal{P}^+$ the set of dominant integral weights $\{\lambda \in \mathcal{P} \mid \forall i \in I, \lambda(h_i) \in \mathbb{N}\}$. For each $i \in I$, $\Lambda_i \in \mathcal{P}^+$ is a dominant integral weight determined modulo the subgroup $\{\lambda \in \mathcal{P} \mid \forall i \in I, \lambda(h_i) = 0\} (\subseteq \mathcal{P})$ by the condition $\Lambda_i(h_j) = \delta_{ij}$ for all $j \in I$.

Recall that the Weyl group $W = W(A)$ is a subgroup of $\mathrm{GL}(\mathfrak{h}^*)$ generated by $\{s_i : \mathfrak{h}^* \xrightarrow{\sim} \mathfrak{h}^*, \lambda \longmapsto \lambda - \lambda(h_i)\alpha_i \mid i \in I\}$ and that W also acts on $\mathcal{P} (\subseteq \mathfrak{h}^*)$.

In the following, we use the usual abberiviation such as $v_i = v^{d_i}$, $[n]_\ell = \sum_{k=1}^{n} v^{(n+1-2k)\ell}$, $[n]_\ell! = \prod_{m=1}^{n} [m]_\ell$ and $\begin{bmatrix} n \\ m \end{bmatrix}_i = \frac{[n]_i!}{[m]_i![n-m]_i!}$ for $i \in I$ and $n \geq m \geq 0$.

Definition 4.1. *The quantum group $U_v = U_v(A)$ is an associative $\mathbf{k}$-algebra with 1 generated by $\{e_i, f_i \mid i \in I\} \cup \{v^h \mid h \in \mathcal{P}^\vee\}$ with the following definition relations.*

(a) $v^0 = 1$ *and* $v^h v^{h'} = v^{h+h'}$ *for any* $h, h' \in \mathcal{P}$,

(b) $v^{-h} e_i v^h = v^{\alpha_i(h)} e_i, v^{-h} f_i v^h = v^{-\alpha_i(h)} f_i$ *for any* $i \in I$ *and* $h \in \mathcal{P}$,

(c) $e_i f_j - f_j e_i = \delta_{ij}(K_i - K_i^{-1})/(v_i - v_i^{-1})$ *for any* $i, j \in I$,

(d) $\sum_{k=0}^{1-a_{ij}} (-1)^k e_i^{(k)} e_j e_i^{(1-a_{ij}-k)} = 0$ *for any* $i, j \in I$ *with* $i \neq j$,

(e) $\sum_{k=0}^{1-a_{ij}} (-1)^k f_i^{(k)} f_j f_i^{(1-a_{ij}-k)} = 0$ *for any* $i, j \in I$ *with* $i \neq j$,

where $K_i = v^{d_i h_i}, v_i = v^{d_i}$ *and* $e_i^{(n)} = e_i^n/[n]_{d_i}!, f_i^{(n)} = f_i^n/[n]_{d_i}!$.

Remark 4.2. U_v has $\mathbf{k}$-antiinvolution σ and $\mathbb{Q}$-antiinvolution Ω, Υ defined by

$$\begin{aligned} &\sigma(e_i) = f_i, \quad \sigma(f_i) = e_i, \quad \sigma(v^h) = v^h, \\ &\Omega(e_i) = f_i, \quad \Omega(f_i) = e_i, \quad \Omega(v^h) = v^{-h}, \quad \Omega(v) = v^{-1}, \\ &\Upsilon(e_i) = v_i f_i K_i^{-1}, \quad \Upsilon(f_i) = v_i^{-1} K_i e_i, \quad \Upsilon(v^h) = v^{-h}, \quad \Upsilon(v) = v^{-1}. \end{aligned}$$

Proposition 4.3 ([Lus, §3.2])**.** *Define the three $\mathbf{k}$-subalgebras $U_v^{\pm}, U_v^0$ of U_v by*

$$U_v^+ = \langle e_i \mid i \in I \rangle, \quad U_v^- = \langle f_i \mid i \in I \rangle, \quad U_v^0 = \langle v^h \mid h \in \mathcal{P} \rangle.$$

(a) *the canonical map* $U_v^- \otimes_{\mathbf{k}} U_v^0 \otimes_{\mathbf{k}} U_v^+ \to U_v$ *is a $\mathbf{k}$-vector space isomorphism,*

(b) U_v^0 *is canonically isomorphic to the group $\mathbf{k}$-algebra* $\mathbf{k}[\mathcal{P}]$.

Definition 4.4. *We define the following two maps both are assured by Proposition 4.3.*

(a) *the Harish-Chandra projection* $\mathsf{HC} : U_v \to U_v^0$ *which is a $\mathbf{k}$-linear projector from* $U_v = U_v^0 \oplus ((\sum_{i \in I} f_i U_v) + (\sum_{i \in I} U_v e_i))$ *to* U_v^0.

(b) *the evaluation map* $\mathsf{ev}_\lambda : U_v^0 \to \mathbf{k}$ *which is an $\mathbf{k}$-algebra homomorphism determined by the assignment* $\mathsf{ev}_\lambda(v^h) = v^{\lambda(h)}$ *for each* $h \in \mathcal{P}$.

4.2. Shapovalov forms.

Definition 4.5 ([Lus, §3.4.5])**.** *For $\lambda \in \mathcal{P}$, the Verma module is a left U_v-module*

$$M(\lambda) = U_v / (\sum_{h \in \mathcal{P}} U_v (v^h - v^{\lambda(h)}) + \sum_{i \in I} U_v e_i).$$

Remark 4.6. In virtue of Proposition 4.3, as a U_v^--module $M(\lambda)$ is free of rank 1 and we see that $M(\lambda)$ has a proper maximum U_v-submodule $K(\lambda)$. We reserve the symbol φ_λ for the U_v^--linear isomorphism $\varphi_\lambda : U_v^- \xrightarrow{\sim} M(\lambda), u \mapsto u + \sum_{h \in \mathcal{P}} U_v(v^h - v^{\lambda(h)}) + \sum_{i \in I} U_v e_i$.

Definition 4.7. *For $\lambda \in \mathcal{P}$, we define an irreducible U_v-module $V(\lambda) = M(\lambda)/K(\lambda)$.*

Proposition 4.8. *Let us take $\lambda \in \mathcal{P}$.*

(a) *There exists a unique non-degenerate symmetric* $\mathbf{k}$*-bilinear form* $\langle , \rangle_{\mathsf{Sh}} : V(\lambda)\times V(\lambda) \to \mathbf{k}$ *(called the Shapovalov form) such that* $\langle 1_\lambda, 1_\lambda\rangle_{\mathsf{Sh}} = 1$ *and* $\langle uv, w\rangle_{\mathsf{Sh}} = \langle v, \sigma(u)w\rangle_{\mathsf{Sh}}$ *for all* $v, w \in V(\lambda)$ *and* $u \in U_v$*,*

(b) *There exist unique bi-additive non-degenerate maps* $\langle , \rangle_{\mathsf{QSh}} : V(\lambda) \times V(\lambda) \to \mathbf{k}$ *and* $\langle , \rangle_{\mathsf{RSh}} : V(\lambda) \times V(\lambda) \to \mathbf{k}$ *such that for all* $X \in \{\mathsf{QSh}, \mathsf{RSh}\}$
  (i) $\langle aw_1, w_2\rangle_X = \sigma(a)\langle w_1, w_2\rangle_X$*,* $\langle w_1, aw_2\rangle_X = a\langle w_1, w_2\rangle_X$ *and* $\langle w_1, w_2\rangle_X = \sigma(\langle w_2, w_1\rangle_X)$*,*
  (ii) $\langle 1_\lambda, 1_\lambda\rangle_X = 1$ *and* $\langle uw_1, w_2\rangle_{\mathsf{QSh}} = \langle w_1, \Omega(u)w_2\rangle_{\mathsf{QSh}}$*,* $\langle uw_1, w_2\rangle_{\mathsf{QSh}} = \langle w_1, \Upsilon(u)w_2\rangle_{\mathsf{RSh}}$*.*

for all $w_1, w_2 \in V(\lambda)$*,* $u \in U_v$ *and* $a \in \mathbf{k}$*.*

Proof. All cases are similar and standard. Let $X \in \{\mathsf{Sh}, \mathsf{QSh}, \mathsf{RSh}\}$ and put $\Xi = \sigma, \Omega, \Upsilon$ according to $X = \mathsf{Sh}, \mathsf{QSh}, \mathsf{RSh}$ respectively. We just sketch the constructions of $\langle , \rangle_X$ since they are needed in the proof of Proposition 4.16.

Define $\langle , \rangle'_X : M(\lambda) \times M(\lambda) \to \mathbf{k}$ by $\langle w_1, w_2\rangle'_X = \mathsf{ev}_\lambda(\mathsf{HC}(\Xi(\varphi_\lambda^{-1}(w_1))\varphi_\lambda^{-1}(w_2)))$. Then, we see that $\mathsf{Rad}(\langle , \rangle'_X) = K(\lambda)$ and $\langle , \rangle'_X$ decent to $\langle , \rangle_X : V(\lambda) \times V(\lambda) \to \mathbf{k}$ which satisfy the desired conditions. ☐

4.3. **Lusztig lattices.**

Theorem 4.9 ([Lus, Theorem 14.4.3],[Lus2, Theorem 4.5])**.** *Let* $U_v^{\mathscr{A}}$ *be an* $\mathscr{A}$*-subalgebra generated by* $\{e_i^{(n)}, f_i^{(n)}, K_i^{\pm 1} \mid i \in I, n \geq 0\}$*. Then,* $U_v^{\mathscr{A}}$ *is an* $\mathscr{A}$*-lattice of* U_v*.*

Definition 4.10 ([Lus, §3.5])**.** *A* U_v*-module* M *is called integrable*

(i) $M = \oplus_{\nu\in\mathfrak{h}^*} M_\nu$ *with* $\dim M_\nu < +\infty$ *where* $M_\nu = \{m \in M \mid \forall h \in \mathcal{P}^\vee, v^h m = v^{\nu(h)}m\}$*,*
(ii) *for any* $m \in M$ *and* $i \in I$*, there exists some* $n > 0$ *such that* $f_i^n m = e_i^n m = 0$*.*

Remark 4.11 ([Lus, Proposition 3.5.8, Proposition 5.2.7])**.** It is well-known that
(a) for $\lambda \in \mathcal{P}$, $V(\lambda)$ is integrable if and only if $\lambda \in \mathcal{P}^+$.
(b) for $\lambda \in \mathcal{P}^+$, the set of weights $P(\lambda) := \{\mu \in \mathfrak{h}^* \mid V(\lambda)_\mu \neq 0\}$ of $V(\lambda)$ is W-invariant.

Theorem 4.12 ([Lus, Theorem 14.4.11])**.** *Assume* $\lambda \in \mathcal{P}^+$*. Then,* $V(\lambda)^{\mathscr{A}} := U_v^{\mathscr{A}} 1_\lambda$ *is an* $\mathscr{A}$*-lattice of* $V(\lambda)$ *compatible with the weight space decomposition of* $V(\lambda)$*. In other words,* $V(\lambda)_\nu^{\mathscr{A}} := V(\lambda)_\nu \cap V(\lambda)^{\mathscr{A}}$ *is an* $\mathscr{A}$*-lattice of* $V(\lambda)_\nu$ *for all* $\nu \in P(\lambda)$ *and we have* $V(\lambda)^{\mathscr{A}} = \oplus_{\nu\in P(\lambda)} V(\lambda)_\nu^{\mathscr{A}}$*.*

Definition 4.13. *For* $\lambda \in \mathcal{P}^+$ *and* $\mu \in P(\lambda)$*, we define* $\mathsf{Sh}_{\lambda,\mu} = \det \mathsf{Sh}^{\mathsf{M}}_{\lambda,\mu} \in \mathbf{k}^\times/v^{2\mathbb{Z}}$ *and*

$$\begin{aligned}\mathsf{Sh}^{\mathsf{M}}_{\lambda,\mu} &= \mathsf{Sh}^{\mathsf{M}}_{\lambda,\mu}(A) = \mathsf{GM}_{\mathbf{k},\mathsf{id}_{\mathbf{k}},\mathscr{A}}(V(\lambda)_\mu, \mathsf{Sh}, V(\lambda)_\mu^{\mathscr{A}}),\\ \mathsf{QSh}^{\mathsf{M}}_{\lambda,\mu} &= \mathsf{QSh}^{\mathsf{M}}_{\lambda,\mu}(A) = \mathsf{GM}_{\mathbf{k},\tau,\mathscr{A}}(V(\lambda)_\mu, \mathsf{QSh}, V(\lambda)_\mu^{\mathscr{A}}),\\ \mathsf{RSh}^{\mathsf{M}}_{\lambda,\mu} &= \mathsf{RSh}^{\mathsf{M}}_{\lambda,\mu}(A) = \mathsf{GM}_{\mathbf{k},\tau,\mathscr{A}}(V(\lambda)_\mu, \mathsf{RSh}, V(\lambda)_\mu^{\mathscr{A}}).\end{aligned}$$

Remark 4.14. Theorem 4.9 and Theorem 4.12 allow us to specialize the integrable highest weight modules. For any $\xi \in \mathbb{C}^\times$ and $\lambda \in \mathcal{P}^+$, the specialized module $V(\lambda)|_{v=\xi} := \mathbb{C}_\xi \otimes_{\mathscr{A}} V(\lambda)^{\mathscr{A}}$ is a module over $U_v|_{v=\xi} := \mathbb{C}_\xi \otimes_{\mathscr{A}} U_v^{\mathscr{A}}$ where $\mathbb{C}_\xi = \mathbb{C}$ is an $\mathscr{A}$-module where v acts as the multiplication of ξ.

Remark 4.15. For $\lambda \in \mathcal{P}^+$ and $\xi \in \mathbb{C}^\times$, $V(\lambda)|_{v=\xi}$ is irreducible if and only if $\mathsf{Sh}_{\lambda,\mu}(\xi) \neq 0$ for all $\mu \in P(\lambda)$. Thus, for a generic $\xi \in \mathbb{C}^\times$, $V(\lambda)|_{v=\xi}$ is irreducible. It is expected that $V(\lambda)|_{v=\xi}$ is irreducible when ξ is not a root of unity [Kas, Problem 2].

Proposition 4.16. *Let* $\lambda \in \mathcal{P}^+$ *and take* $\mu \in P(\lambda)$*. We may assume* $\mathsf{Sh}^{\mathsf{M}}_{\lambda,\mu} = \mathsf{QSh}^{\mathsf{M}}_{\lambda,\mu}$ *and* $D\mathsf{QSh}^{M}_{\lambda,\mu} = \mathsf{RSh}^{M}_{\lambda,\mu}$ *for some diagonal matrix* D *all of whose diagonal entries belong to* $v^{\mathbb{Z}}$*.*

Proof. By the constructions mentioned in the proof of Proposition 4.8, it is enough to show that we can choose an $\mathscr{A}$-basis of $V(\lambda)^{\mathscr{A}}_{\mu}$ as a subset of $\{f^{(n_1)}_{i_1}\cdots f^{(n_s)}_{i_s}1_\lambda \mid {}^{s\geq 0}_{1\leq\forall j\leq s, n_j\geq 1, i_j\in I}\}$. In virtue of [Lak, Theorem 6.5], it is possible[1]. $\square$

Corollary 4.17. *For $\lambda\in\mathcal{P}^+$ and $\mu\in P(\lambda)$, we have* $\det\mathsf{QSh}^M_{\lambda,\mu}=\det\mathsf{RSh}^M_{\lambda,\mu}$ *in* $\mathbf{k}^\times$.

Proof. We may assume $D\mathsf{QSh}^M_{\lambda,\mu}=\mathsf{RSh}^M_{\lambda,\mu}$ for some D with $\det D\in v^{\mathbb{Z}}$. Since ${}^{\mathrm{tr}}\mathsf{QSh}^M_{\lambda,\mu}=\tau(\mathsf{QSh}^M_{\lambda,\mu})$ and ${}^{\mathrm{tr}}\mathsf{RSh}^M_{\lambda,\mu}=\tau(\mathsf{RSh}^M_{\lambda,\mu})$, we easily see that $\det D=\tau(\det D)$. $\square$

Proposition 4.18. *For $\lambda\in\mathcal{P}^+$, $i\in I$ and $\mu\in P(\lambda)$, we may assume* $\mathsf{QSh}^{\mathsf{M}}_{\lambda,\mu}=\mathsf{QSh}^{\mathsf{M}}_{\lambda,s_i(\mu)}$.

Proof. Since $V(\lambda)$ is integrable, we have for each $i\in I$ mutually inverse linear operators $T'_{i,-1}=\sum_{\nu\in P(\lambda)}T'^{(\nu)}_{i,-1}$ and $T''_{i,+1}=\sum_{\nu\in P(\lambda)}T''^{(\nu)}_{i,+1}$ on $V(\lambda)$ where (see [Lus, §5.2])

$$T'^{(\nu)}_{i,-1}:V(\lambda)_\nu\xrightarrow{\sim}V(\lambda)_{s_i(\nu)},\quad w\longmapsto\sum_{\substack{a,b,c\geq 0\\ a-b+c=\nu(h_i)}}(-1)^b v_i^{ac-b}f_i^{(a)}e_i^{(b)}f_i^{(c)}w,$$

$$T''^{(\nu)}_{i,+1}:V(\lambda)_\nu\xrightarrow{\sim}V(\lambda)_{s_i(\nu)},\quad w\longmapsto\sum_{\substack{a,b,c\geq 0\\ -a+b-c=\nu(h_i)}}(-1)^b v_i^{b-ac}e_i^{(a)}f_i^{(b)}e_i^{(c)}w.$$

It is clear that we have $\langle T'_{i,-1}(v),T'_{i,-1}(w)\rangle_{\mathsf{QSh}}=\langle v,T''_{i,+1}(T'_{i,-1}(w))\rangle_{\mathsf{QSh}}=\langle v,w\rangle_{\mathsf{QSh}}$. Since $T'_{i,-1}$ (and $T''_{i,+1}$) preserves $V(\lambda)^{\mathscr{A}}$ [Lus, Proposition 41.2.2], we are done. $\square$

5. SHAPOVALOV DETERMINANTS OF THE BASIC REPRESENTATIONS

5.1. Untwisted affine A,D,E case.

Let $X=(a_{ij})_{i,j\in I}$ be a Cartan matrix of type A,D,E and let $\widehat{X}=X^{(1)}$ be the extended Cartan matrix of X indexed by $\widehat{I}=\{0\}\sqcup I$ as in Figure 1. Let $(a_i)_{i\in\widehat{I}}$ be the numerical labels of $\widehat{X}$ in Figure 1 and let $\delta=\sum_{i\in\widehat{I}}a_i\alpha_i$.

Remark 5.1 ([Kac, §12.6])**.** Let $W=W(\widehat{X})$. We have

$$P(\Lambda_0)=\{w\Lambda_0-d\delta\mid w\in W,d\geq 0\}(=\{w\Lambda_0-d\delta\mid w\in W/W_0,d\geq 0\})\tag{5.1}$$

Note that $W/W_0\cong W(X)$ where $W_0=\{w\in W\mid w\Lambda_0=\Lambda_0\}$.

We shall very briefly review the vertex operator construction of $V(\Lambda_0)$, i.e., an explicit realization of $V(\Lambda_0)$ as the tensor product of a polynomial algebra and a (twisted) group algebra [FJ] (however, we will mainly refer [CJ] instead). Recall the Drinfeld new realization of $U_v(\widehat{X})$ which gives loop-like generators [Dri].

Theorem 5.2. *As a* $\mathbf{k}$*-algebra,* $U_v(\widehat{X})$ *is isomorphic to the* $\mathbf{k}$*-algebra generated by* $\{x^\pm_{i,s}\mid i\in I,s\in\mathbb{Z}\}\cup\{h_{i,r}\mid i\in I,r\in\mathbb{Z}\setminus\{0\}\}\cup\{C^{\pm 1},D^{\pm 1},K_i^{\pm 1}\mid i\in I\}$ *with the following defining relations.*

(a) $C^{\pm 1}$ *are central and* $CC^{-1}=C^{-1}C=DD^{-1}=D^{-1}D=K_iK_i^{-1}=K_i^{-1}K_i=1$,

(b) $K_iK_j=K_jK_i$, $K_ih_{j,r}=h_{j,r}K_i$, $K_ix^\pm_{j,s}K_i^{-1}=v^{\pm a_{ij}}x^\pm_{j,s}$,

(c) $DK_i=K_iD$, $Dh_{j,r}D^{-1}=v^rh_{j,r}$, $Dx^\pm_{j,s}D^{-1}=v^rx^\pm_{j,s}$,

---

[1] For the basic representations of $U_v(A^{(1)}_n),U_v(A^{(2)}_{2n-1}),U_v(A^{(2)}_{2n}),U_v(D^{(1)}_n),U_v(D^{(2)}_{n+1}),U_v(B^{(1)}_n)$, we can choose such monomial $\mathscr{A}$-bases using combinatorial algorithms [LLT, LT, KKw].


A_1
α_1
A_ℓ
α_1 α_2 $\alpha_{\ell-1}$ α_ℓ
D_ℓ
$\alpha_{\ell-1}$
α_1 α_2 $\alpha_{\ell-2}$ $\alpha_{\ell-1}$
E_6
α_6
α_1 α_2 α_3 α_4 α_5
E_7
α_7
α_1 α_2 α_3 α_4 α_5 α_6
E_8
α_8
α_1 α_2 α_3 α_4 α_5 α_6 α_7
$A_1^{(1)}$
1 1
$A_\ell^{(1)}$
1
1 1 1
$D_\ell^{(1)}$
1 1
1 2 2 2 1
$E_6^{(1)}$
1
2
1 2 3 2 1
$E_7^{(1)}$
2
1 2 3 4 3 2 1
$E_8^{(1)}$
3
1 2 3 4 5 6 4 2

FIGURE 1. Finite and untwisted affine Dynkin diagrams of A,D,E.

(d) $[h_{i,r}, h_{j,r'}] = \delta_{r+r',0}\frac{[ra_{ij}]}{r}\frac{C^r - C^{-r}}{v-v^{-1}}$, $[x^+_{i,s}, x^-_{j,s'}] = \delta_{i,j}\frac{C^{-s'}\psi^+_{i,s+s'} - C^{-s}\psi^-_{i,s+s'}}{v-v^{-1}}$,

(e) $[h_{i,\pm r}, x^\pm_{j,s}] = \pm\frac{[ra_{ij}]}{r}x^\pm_{j,s\pm r}$, $[h_{i,\mp r}, x^\pm_{j,s}] = \pm\frac{[ra_{ij}]}{r}C^r x^\pm_{j,s\pm r}$,

(f) $x^\pm_{i,s+1}x^\pm_{j,s'} - v^{\pm a_{ij}}x^\pm_{j,s'}x^\pm_{i,s+1} = v^{\pm a_{ij}}x^\pm_{i,s}x^\pm_{j,s'+1} - x^\pm_{j,s'+1}x^\pm_{i,s}$,

(g) $\sum_{\pi\in\mathfrak{S}_{1-a_{pq}}}\sum_{\substack{\boldsymbol{r}=(r_\ell)\in\mathbb{Z}^{1-a_{pq}}\\ 0\le k\le 1-a_{pq}}}(-1)^k\begin{bmatrix}1-a_{pq}\\ k\end{bmatrix}x^\pm_{p,r_{\pi(1)}}\cdots x^\pm_{p,r_{\pi(k)}}x^\pm_{q,s}x^\pm_{p,r_{\pi(k+1)}}\cdots x^\pm_{p,r_{\pi(m)}} = 0$,

for all $i,j,p,q \in I$ *with* $p \neq q$ *and* $r, r' \in \mathbb{Z}\setminus\{0\}$, $s, s' \in \mathbb{Z}$ *where we set* $\psi^+_{-s} = \psi^-_s = 0$ *for* $s \geq 1$ *and* $\sum_{t\geq 0}\psi^\pm_{i,\pm t}z^{\pm t} = K_i^{\pm 1}\exp(\pm(v-v^{-1})\sum_{r\geq 1}h_{i,\pm r}z^{\pm r})$.

Remark 5.3. Actually, as shown by Beck [Bec] (see also [BCP, Theorem 1]), we may choose the generators in Theorem 5.2 as particular elements in $U_v(\widehat{X})$, some of them are constructed utilizing Lusztig's braid group action on $U_v(\widehat{X})$ from the Chevalley generators of $U_v(\widehat{X})$. Although we omit the construction, in the rest of this paper we assume this special choice for each generator in Theorem 5.2. The choice is used essentially in that

(a) an $\mathscr{A}$-subalgebra generated by $\{(x^\pm_{i,n})^r/[r]! \mid i \in I, r \geq 0\}$ coincides with $U_v^{\pm,\mathscr{A}}$ [BCP, Corollary 2.2] where $U_v^{+,\mathscr{A}} = \langle e_i^{(r)} \mid i \in I, r \geq 0\rangle_{\mathscr{A}\text{-alg}}$ and $U_v^{-,\mathscr{A}} = \langle f_i^{(r)} \mid i \in I, r \geq 0\rangle_{\mathscr{A}\text{-alg}}$. This is used in the proof of Theorem 5.5 (ii),

(b) we have $\Omega(h_{i,\pm r}) = h_{i,\mp r}$ for $i \in I$ and $r \geq 1$ [BCP, Lemma 3]. This will be used in the proof of Theorem 5.6.

Remark 5.4. Let $U^-(0)$ be the $\mathbf{k}$-subalgebra of $U_v(\widehat{X})$ generated by $\{h_{i,-r} \mid i \in I, r \geq 1\}$. Clearly, it is a commutative algebra. In virtue of [BCP, Proposition 1.3], $\{h_{i,-r} \mid i \in I, r \geq 1\}$ is algebraically independent over $\mathbf{k}$.

Theorem 5.5 ([CJ, Theorem2, Theorem 3]). *Let* $Q = \oplus_{p\in I}\mathbb{Z}\alpha_p$ *be the root lattice of* X. *We can put a* $U_v(\widehat{X})$*-module structure on a* $\mathbf{k}$*-vector space* $V := U^-(0)\otimes_{\mathbf{k}}\mathbf{k}[Q]$ *by* *extending the following assignment of action $\odot$ on V*

$$h_{i,-s}\odot(h_{i_1,-r_1}\cdots h_{i_j,-r_j}\otimes e^{\beta}) = h_{i,-s}h_{i_1,-r_1}\cdots h_{i_j,-r_j}\otimes e^{\beta},$$

$$h_{i,s}\odot(h_{i_1,-r_1}\cdots h_{i_j,-r_j}\otimes e^{\beta}) = \sum_{k=1}^{j} h_{i_1,-r_1}\cdots h_{i_{k-1},-r_{k-1}}\frac{\delta_{s,r_k}[sa_{i,i_k}][s]}{s}h_{i_{k+1},-r_{k+1}}\cdots h_{i_j,-r_j}\otimes e^{\beta},$$

to all the generators in Theorem 5.2 and can obtain a $U_v(\widehat{X})$-module isomorphism $\varphi : V \xrightarrow{\sim} V(\Lambda_0)$ such that

(i) $\varphi(h_{i_1,-r_1}\cdots h_{i_j,-r_j}\otimes e^{\beta}) \in V(\Lambda_0)_{\Lambda_0-\beta-(\sum_{k=1}^{j} r_k+(\beta,\beta)/2)\delta}$,

(ii) $\varphi(\mathscr{A}[\widetilde{P}^-_{i,r} \mid i\in I, r\geq 1]\otimes_{\mathscr{A}}\mathscr{A}[Q]) = V(\Lambda_0)^{\mathscr{A}}$ *where* $1+\sum_{r\geq 1}\widetilde{P}^-_{i,r}z^r = \exp(\sum_{r\geq 1}\frac{h_{i,-r}}{[r]}z^r)$.

Here $(i_1,\cdots,i_j)\in I^j, (r_1,\cdots,r_j)\in\mathbb{N}_+^j, \beta=\sum_{p\in I}b_p\alpha_p\in Q$, $i\in I$, $s\geq 1$, $(\beta,\beta) = \sum_{p,q\in I}b_pa_{pq}b_q$ *and we regard Q as an additive subgroup of the weight lattice $\mathcal{P}$ of $\widehat{X}$.*

Theorem 5.6. *Let $X=(a_{ij})_{i,j\in I}$ be a Cartan matrix of type A,D,E and let $\widehat{X}$ be the extended Cartan matrix of X indexed by $\widehat{I}=\{0\}\sqcup I$ as in Figure 1. We have*

$$\mathsf{Sh}_{\Lambda_0,w\Lambda_0-d\delta}(\widehat{X}) = \prod_{s=1}^{d}(\det[X]_s)^{\sum_{\lambda\in\mathsf{Par}(d)}\frac{m_s(\lambda)}{|I|}N_\lambda}$$

in $\mathbf{k}^\times/v^{2\mathbb{Z}}$ *for all* $w\in W$ *and* $d\geq 0$ *where* $N_\lambda=\prod_{u\geq 1}\binom{m_u(\lambda)+|I|-1}{m_u(\lambda)}$ *and* $[X]_s=([a_{ij}]_s)_{i,j\in I}$.

Proof. By Proposition 4.16 and Proposition 4.18, we shall calculate $\det\mathsf{QSh}^{\mathsf{M}}_{\Lambda_0,\Lambda_0-d\delta}(\widehat{X})$ instead of $\mathsf{Sh}_{\Lambda_0,w\Lambda_0-d\delta}(\widehat{X})$. By Theorem 5.5, we see

(i) $V(\Lambda_0)_{\Lambda_0-d\delta}$ has a $\mathbf{k}$-basis $\{h'_{i_1,-r_1}\cdots h'_{i_j,-r_j}\otimes e^0 \mid (i_\ell)\in I^j, (r_\ell)\in\mathbb{N}_+^j \text{ with } \sum_{k=1}^{j}r_k=d\}$,

(ii) $V(\Lambda_0)^{\mathscr{A}}_{\Lambda_0-d\delta}$ has an $\mathscr{A}$-basis $\{\widetilde{P}^-_{i_1,r_1}\cdots\widetilde{P}^-_{i_j,r_j}\otimes e^0 \mid (i_\ell)\in I^j, (r_\ell)\in\mathbb{N}_+^j \text{ with } \sum_{k=1}^{j}r_k=d\}$,

where $h'_{i,\pm r}=h_{i,\pm r}/[r]$ for $i\in I$ and $r\geq 1$. Note that $1+\sum_{r\geq 1}\widetilde{P}^-_{i,r}z^r=\exp(\sum_{r\geq 1}h'_{i,-r}z^r)$.

Since $\Omega(h'_{i,-r})=h'_{i,r}$ (see Remark 5.3) and

$$(sh'_{i,s})\odot(h'_{i_1,-r_1}\cdots h'_{i_j,-r_j}\otimes e^0) = \sum_{k=1}^{j}\delta_{s,r_k}[a_{i,i_k}]_s h'_{i_1,-r_1}\cdots h'_{i_{k-1},-r_{k-1}}h'_{i_{k+1},-r_{k+1}}\cdots h'_{i_j,-r_j}\otimes e^0,$$

we can apply Corollary 3.8 and get $\det\mathsf{QSh}^{\mathsf{M}}_{\Lambda_0,\Lambda_0-d\delta}(A)=\prod_{s=1}^{d}(\det[A]_s)^{\sum_{\lambda\in\mathsf{Par}(d)}\frac{m_s(\lambda)}{|I|}N_\lambda}$. □

Corollary 5.7. *Let $\mathfrak{g}_0$ be a finite-dimensional simple simply-laced complex Lie algebra.*

(a) *If $\xi\in\mathbb{C}^\times$ is not a root of unity, then $V(\Lambda_0)|_{v=\xi}$ is irreducible.*

(b) *Let $\mathfrak{g}_0=\mathfrak{g}(X)$ and let $\ell\geq 1$. $V(\Lambda_0)|_{v=\exp(\frac{2\pi\sqrt{-1}}{\ell})}$ is irreducible if and only if*

  (i) $\mathsf{gcd}(\ell,2n)=1,2$ *when* $X=A_{n-1}$ *for* $n\geq 2$,

  (ii) $\ell\notin 4\mathbb{Z}$ *(resp.* $\notin 3\mathbb{Z},\notin 4\mathbb{Z},\notin 60\mathbb{Z}$*) when* $X=D_n$ *for* $n\geq 4$ *(resp.* E_6,E_7,E_8*).*

Proof. The specialized module is irreducible if and only if $(\det[X]_k)|_{v=\xi}\neq 0$ for all $k\geq 1$. Type by type, $\det([X]_1)$ is given by $\det([A_{n-1}]_1)=[n]$, $\det([D_m]_1)=v^{-m}\Phi_4(v)\Phi_4(v^{m-1})$, $\det([E_6]_1)=v^{-6}\Phi_3(v^2)\Phi_{24}(v)$, $\det([E_7]_1)=v^{-7}\Phi_4(v)\Phi_{36}(v)$, $\det([E_8]_1)=v^{-8}\Phi_{60}(v)$ for $n\geq 2$, $m\geq 4$ where $\Phi_j(x)\in\mathbb{Z}[x]$ is the j-th cyclotomic polynomial. □



Figure 2. Twisted affine Dynkin diagrams of A,D,E.

Remark 5.8. Let $\mathbf{k}$ be a field of characteristic $p \geq 2$. DeConcini-Kac-Kazhdan proved that the modulo-p reduction $\mathbf{k} \otimes_{\mathbb{Z}} V(\Lambda_0)^{\mathbb{Z}}$ of the basic $\widehat{\mathfrak{g}}_0$-module $V(\Lambda_0)$ where $V(\Lambda_0)^{\mathbb{Z}}$ is the Kostant $\mathbb{Z}$-form of $V(\Lambda_0)$ remains irreducible if and only if $\det X \neq 0$ in $\mathbf{k}$ [DcKK, Corollary 3.1]. Corollary 5.7 can be seen a quantum analog of DeConcini-Kac-Kazhdan's result. Note that when $X = E_8$ the modulo-p reduction preserves irreduciblity for any prime $p \geq 2$ because $\det E_8 = 1$. Since the quantum characteristic $\min\{k \geq 1 \mid [k]|_{v=\xi} = 0\}$ takes arbitrary value ≥ 2, quantum case is more subtle than the classical case.

5.2. **Twisted affine A,D,E case.** Let $\widehat{X} = (a_{ij})_{i,j\in\widehat{I}=\{\varepsilon\}\sqcup I}$ be a twisted affine GCM as in Figure 2 where $\varepsilon = \ell$ if $\widehat{X} = A_{2\ell}^{(2)}$ and otherwise $\varepsilon = 0$. Let $\delta = \sum_{i\in\widehat{I}} a_i\alpha_i$ where $(a_i)_{i\in\widehat{I}}$ are the numerical labels of $\widehat{X}$. As in untwisted cases (see Remark 5.1), it is known that $P(\Lambda_\varepsilon) = \{w\Lambda_\varepsilon - d\delta \mid w \in W(\widehat{X}), d \geq 0\}$ [Kac, §12.6].

We discuss on $\mathsf{Sh}_{\Lambda_\varepsilon, w\Lambda_\varepsilon - d\delta}(\widehat{X})$ and give a conjectural formula for it. Our motivation comes from the fact that the Cartan determinant of a faithful p-block of the Schur double covers of the symmetric groups coincides with $\mathsf{Sh}_{\Lambda_\ell, w\Lambda_\ell - d\delta}(A_{p-1}^{(2)})|_{v=1}$ for suitable $w \in W(A_{p-1}^{(2)})$ and $d \geq 0$ when $p = 2\ell + 1$ is an odd prime [BO2, Theorem 5.6].

Definition 5.9 ([BKM, §2.2]). *We define* $\{n\}_s = \frac{v^{ns}+(-1)^s v^{-ns}}{v^s+(-1)^s v^{-s}} (\in \mathscr{A})$ *for* $n, s \geq 1$ *with* $n \notin 2\mathbb{Z}$. *Note that* $\{n\}_s|_{v=1} = n$ *if* s *is odd, but* $\{n\}_s|_{v=1} = 1$ *if* s *is even.*

Recall Theorem 5.6 stating that for $p \geq 2$, we have $\mathsf{Sh}_{\Lambda_0, w\Lambda_0 - d\delta}(A_{p-1}^{(1)}) = \prod_{s=1}^{d} [p]_s^{N_{d,s,p}}$ for all $w \in W(A_{p-1}^{(1)})$ and $d \geq 0$ where $N_{p,d,s} = \sum_{\lambda\in\mathsf{Par}(d)} \frac{m_s(\lambda)}{p-1} \prod_{u\geq 1} \binom{m_u(\lambda)+p-2}{m_u(\lambda)}$.

Conjecture 5.10. Let $p \geq 3$ be an odd integer and put $\ell = (p-1)/2$. We have $\mathsf{Sh}_{\Lambda_\ell, w\Lambda_\ell - d\delta}(A_{p-1}^{(2)}) = \prod_{s=1}^{d} \{p\}_s^{N_{\ell,d,s}}$ for all $w \in W(A_{p-1}^{(2)})$ and $d \geq 0$.

Remark 5.11. Conjecture 5.10 should be proven similarly as Theorem 5.6 if we have

(a) a proof of Drinfeld new realization of $U_v(A_{p-1}^{(2)})$ (see [Dri]),
(b) a proof of vertex operator construction of $U_v(A_{p-1}^{(2)})$-module $V(\Lambda_{(p-1)/2})$ (see [Jin, JM]),
(c) a good choice of Drinfeld loop-like generators which affords us the Lusztig lattice in terms of Drinfeld generators (a candidate is the PBW generators in [BeNa]).

However, (a) is not achieved so far in general (thus, logically (b) nor (c) neither). See also the introduction of [Her].

Remark 5.12. In virtue of [Aka, Appendix B] and [Jin, JM], we can prove Conjecture 5.10 when $p = 3$ by a similar argument of [CJ, Theorem 3] and Theorem 5.6.

Conjecture 5.13. For a twisted affine A,D,E diagram $X = X_N^{(r)}$ (where $r \geq 2$), we assign the quantities $n, k \in \mathbb{N}_+$ and $\alpha, \beta \in \mathscr{A}$ as follows.

$X_N^{(r)}$	$A_{2n-1}^{(2)} (n \geq 3)$	$A_{2n}^{(2)} (n \geq 1)$	$D_{n+1}^{(2)} (n \geq 1)$	$E_6^{(2)}$	$D_4^{(3)}$
n	n	n	n	4	2
k	$n-1$	n	1	2	1
α	$[2]_n$	$[2n+1]^{\mathsf{su}}$	$[2]_n$	$\{3\}_2$	$[3]^{\mathsf{su}}$
β	$[n]$	$[2n+1]$	$[2]$	$[3]$	$[2]$

Here for an odd integer $p \geq 1$, we define $[p]^{\mathsf{su}} = (v^p + v^{-p})/(v + v^{-1})$. Note that for an even integer $m \geq 2$, we have $\{p\}_m = [p]^{\mathsf{su}}|_{v=v^m}$. Put

$$\gamma_{X,s} = \begin{cases} \alpha|_{v=v^s} & (s \in r\mathbb{Z}) \\ \beta|_{v=v^s} & (s \notin r\mathbb{Z}), \end{cases} \qquad f_{X,s} = \begin{cases} n & (s \in r\mathbb{Z}) \\ k & (s \notin r\mathbb{Z}), \end{cases}$$

then, for all $w \in W(\widehat{X})$ and $d \geq 0$ we have $\mathsf{Sh}_{\Lambda_\varepsilon, w\Lambda_\varepsilon - d\delta}(\widehat{X}) = \prod_{s=1}^{d} \gamma_{X,s}^{N_{X,d,s}}$ where

$$N_{X,d,s} = \sum_{\lambda \in \mathsf{Par}(d)} \frac{m_s(\lambda)}{f_{X,s}} \prod_{i \geq 1} \binom{f_{X,i} - 1 + m_i(\lambda)}{m_i(\lambda)}.$$

Remark 5.14. Assume $X_N^{(r)} = (B_{ij})_{i,j \in \widehat{I}} \neq A_{2n}^{(2)}$ for simplicity. Let $X_N = (A_{i,j})_{i,j \in J}$ with $|J| = N$ and let $\mu : X_N \xrightarrow{\sim} X_N$ be a Dynkin diagram automorphism of order r such that the folding procedure to (X_N, μ) gives $X_N^{(r)}$ (See [Kac, §7.9,§8]). In the procedure, I is identified with the set $\{\{\mu^k(j) \mid k \in \mathbb{Z}/r\mathbb{Z}\} \mid j \in J\}$ of μ-orbits of J. We take a map $s : I \to J$ such that i corresponds to $\{\mu^k(s(i)) \mid k \in \mathbb{Z}/r\mathbb{Z}\}$ for all $i \in I$.

For $i \in I$, we put $d_i = a_i^\vee / a_i$ where $a_i^\vee$ be the numerical label of the Langlands dual of $X_N^{(r)}$. Note that we have $d_i = r/|\{\mu^k(s(i)) \mid k \in \mathbb{Z}/r\mathbb{Z}\}|$ for $i \in I$. For $t \geq 1$, we define $I(t) = \{i \in I \mid t \in d_i\mathbb{Z}\}$. Then, the quantities in the above table are designed to satisfy (see also [BK4, §5]) $f_{X,t} = |I(t)|$ and $\det Y^{(t)} = \gamma_{X,t}$ where $Y^{(t)} = (Y_{ij}^{(t)})_{i,j \in I(t)}$ and

$$Y_{ij}^{(t)} = \frac{\sum_{k \in \mathbb{Z}/r\mathbb{Z}} \exp(\frac{2\pi\sqrt{-1}}{r} tk)[A_{s(i),\mu^k(s(j))}]_t}{d_i} = \begin{cases} [B_{ij}]_t & (i = j) \\ B_{ij} & (i \neq j). \end{cases}$$

Remark 5.15. Let $\mathbf{k}$ be a field of characteristic $p \geq 2$ and let $X = X_N^{(r)}$ be a twisted affine A,D,E diagram. Under the assumption that $r \neq 0$ in $\mathbf{k}$, DeConcini-Kac-Kazhdan proved that modulo-p reduction of $\mathfrak{g}(X)$-module $V(\Lambda_\varepsilon)$ remains irreducible if and only if $\det X \neq 0$ in $\mathbf{k}$ [DcKK, Remark 3.1]. Conjecture 5.13 predicts an answer for Kashiwara's problem (see in §1.1) when $\mathfrak{g} = \mathfrak{g}(X)$ and $\lambda(c) = 1$ and it can be seen as a quantum analog of DeConcini-Kac-Kazhdan's result.

6. Graded Cartan determinants of the symmetric groups

Recall that in this paper, we work in the field $\mathbf{k} = \mathbb{Q}(v)$ and its subring $\mathscr{A} = \mathbb{Z}[v, v^{-1}]$.

6.1. Graded representations.

Definition 6.1. *Let A be a finite dimensional algebra over a field $\mathbb{F}$.*

(a) *We denote by $\mathsf{Mod}(A)$ the abelian category consists of finite dimensional left A-modules and A-homomorphisms between them.*

(b) *We define the Cartan matrix* C_A *of* A *to be the matrix* $([\mathsf{PC}(D):D'])_{D,D'\in\mathsf{Irr}(\mathsf{Mod}(A))}\in\mathsf{Mat}_{|\mathsf{Irr}(\mathsf{Mod}(A))|}(\mathbb{Z})$ *where* $\mathsf{PC}(D)$ *is a projective cover of* D.

Definition 6.2. *Let* A *be a graded finite dimensional algebra over a field* $\mathbb{F}$, *i.e.,* A *has a decomposition* $A=\bigoplus_{i\in\mathbb{Z}}A_i$ *into* $\mathbb{F}$*-vector spaces such that* $A_iA_j\subseteq A_{i+j}$ *for all* $i,j\in\mathbb{Z}$.

(a) *We denote by* $\mathsf{Mod}_{\mathsf{gr}}(A)$ *the abelian category consists of finite dimensional left graded* A*-modules and degree preserving* A*-homomorphisms between them.*

(b) *We denote by* $\mathsf{Proj}_{\mathsf{gr}}(A)$ *the abelian category consists of finite dimensional left graded projective* A*-modules and degree preserving* A*-homomorphisms between them.*

(c) *For an object* $M=\bigoplus_{i\in\mathbb{Z}}M_i$ *in* $\mathsf{Mod}_{\mathsf{gr}}(A)$ *and* $k\in\mathbb{Z}$, *we denote by* $M\langle k\rangle$ *the object in* $\mathsf{Mod}_{\mathsf{gr}}(A)$ *such that* $(M\langle k\rangle)_n=M_{k+n}$ *for all* $n\in\mathbb{Z}$.

(d) *For* $k\in\mathbb{Z}$, *the assignment* $(M\xrightarrow{f}N)\longmapsto(M\langle k\rangle\xrightarrow{f}N\langle k\rangle)$ *is obviously an autoequivalence both on* $\mathsf{Mod}_{\mathsf{gr}}(A)$ *and* $\mathsf{Proj}_{\mathsf{gr}}(A)$ *which we denote by* $\langle k\rangle$.

(e) *We define the graded Cartan pairing* $\omega_A:\mathsf{K}_0(\mathsf{Proj}_{\mathsf{gr}}(A))\times\mathsf{K}_0(\mathsf{Mod}_{\mathsf{gr}}(A))\to\mathscr{A}$ *by*

$$\langle[P],[M]\rangle\longmapsto\textstyle\sum_{k\in\mathbb{Z}}\dim_{\mathbb{F}}\mathrm{Hom}_A(P,M\langle k\rangle)v^k.$$

(f) *We define the graded Cartan matrix* C^v_A *of* A *to be the matrix*

$$\left(\textstyle\sum_{k\in\mathbb{Z}}[\mathsf{PC}(D):D'\langle -k\rangle]v^k\right)_{D,D'\in\mathsf{Irr}(\mathsf{Mod}_{\mathsf{gr}}(A))/\sim}\in\mathsf{Mat}_{|\mathsf{Irr}(\mathsf{Mod}_{\mathsf{gr}}(A))/\sim|}(\mathscr{A})$$

where $\mathsf{PC}(D)$ *is a projective cover of* D *in* $\mathsf{Mod}_{\mathsf{gr}}(A)$ *and* $M\sim N$ *if and only if there exists* $k\in\mathbb{Z}$ *such that* $M\langle k\rangle\cong N$ *in* $\mathsf{Mod}_{\mathsf{gr}}(A)$.

Remark 6.3. If $\mathbb{F}$ is a splitting field for A, then $C^v_A={}^{\mathrm{tr}}(\omega_A(\mathsf{PC}(D),\mathsf{PC}(D')))_{D,D'\in\mathsf{Irr}(\mathsf{Mod}_{\mathsf{gr}}(A))/\sim}$.

Remark 6.4. Both $\mathsf{K}_0(\mathsf{Mod}_{\mathsf{gr}}(A))$ and $\mathsf{K}_0(\mathsf{Proj}_{\mathsf{gr}}(A))$ have $\mathscr{A}$-module structures via $v=[\langle -1\rangle]$. ω_A is τ-sesquilinear in that $\omega_A(va,b)=v^{-1}\omega_A(a,b)$ and $\omega_A(a,vb)=v\omega_A(a,b)$ for all $a\in\mathsf{K}_0(\mathsf{Proj}_{\mathsf{gr}}(A))$ and $b\in\mathsf{K}_0(\mathsf{Mod}_{\mathsf{gr}}(A))$.

Remark 6.5. Let C^v_1 and C^v_2 be the graded Cartan matrices of A according to different choice of representatives of $\mathsf{Irr}(\mathsf{Mod}_{\mathsf{gr}}(A))/\sim$. Then, there exists a diagonal matrix D all of whose diagonal entries belong to $v^{\mathbb{Z}}$ such that $C^v_2=\tau({}^{\mathrm{tr}}D)C^v_1D$. Thus, the graded Cartan determinant $\det C^v_A$ is well-defined.

Remark 6.6. It is a routine to prove that the natural map which forgets the grading gives a bijection $\mathsf{Irr}(\mathsf{Mod}_{\mathsf{gr}}(A))/\sim\,\xrightarrow{\sim}\mathsf{Irr}(\mathsf{Mod}(A))$ and $C^v_A|_{v=1}=C_A$.

6.2. **The Khovanov-Lauda-Rouquier algebras.**

Definition 6.7 ([Rou, §3.2.1]). *Let* $\mathbb{F}$ *be a field and let* I *be a finite set. Take a matrix* $Q=(Q_{ij}(u,v))\in\mathsf{Mat}_I(\mathbb{F}[u,v])$ *subject to* $Q_{ii}=0,Q_{ij}(u,v)=Q_{ji}(v,u)$ *for all* $i,j\in I$.

(a) *The KLR algebra* $R_n(\mathbb{F};Q)$ *for* $n\geq 0$ *is an* $\mathbb{F}$*-algebra generated by* $\{x_p,\tau_a,e_\nu\mid 1\leq p\leq n,1\leq a<n,\nu\in I^n\}$ *with the following defining relations for all* $\mu,\nu\in I^n,1\leq p,q\leq n$, $1\leq b<a\leq n-1$.

- $e_\mu e_\nu=\delta_{\mu\nu}e_\mu, 1=\sum_{\mu\in I^n}e_\mu,\ x_px_q=x_qx_p,\ x_pe_\mu=e_\mu x_p,$
- $\tau_a^2e_\nu=Q_{\nu_a,\nu_{a+1}}(x_a,x_{a+1})e_\nu,\ \tau_ae_\mu=e_{s_a(\mu)}\tau_a,$
- $(\tau_ax_{a+1}-x_a\tau_a)e_\nu=(x_{a+1}\tau_a-\tau_ax_a)e_\mu=\delta_{\nu_a,\nu_{a+1}}e_\nu,$
- $(\tau_{b+1}\tau_b\tau_{b+1}-\tau_b\tau_{b+1}\tau_b)e_\nu=\delta_{\nu_b,\nu_{b+2}}((x_{b+2}-x_b)^{-1}(Q_{\nu_b,\nu_{b+1}}(x_{b+2},x_{b+1})-Q_{\nu_b,\nu_{b+1}}(x_b,x_{b+1})))e_\nu.$
- $\tau_a\tau_b=\tau_b\tau_a$ *if* $|a-b|>1,$
- $\tau_ax_p=x_p\tau_a$ *if* $p\neq a,a+1,$

(b) *For* $\beta=\sum_{i\in I}\beta_i\cdot i\in\mathbb{N}[I]$ *with* $n=\mathsf{ht}(\beta):=\sum_{i\in I}\beta_i$, *we define* $R_\beta(\mathbb{F};Q)=R_n(\mathbb{F};Q)e_\beta$ *where* $e_\beta=\sum_{\nu\in\mathsf{Seq}(\beta)}e_\nu$ *and* $\mathsf{Seq}(\beta)=\{(i_j)_{j=1}^n\in I^n\mid\sum_{j=1}^n i_j=\beta\}$.

(c) *For* $\lambda = \sum_{i\in I}\lambda_i\cdot i \in \mathbb{N}[I]$ *and* $\beta\in\mathbb{N}[I]$ *with* $n=\mathsf{ht}(\beta)$*, we define*

$$R^{\lambda}_{n}(\mathbb{F};Q) = R_n(\mathbb{F};Q)/R_n(\mathbb{F};Q)(\textstyle\sum_{\nu\in I^n} x_1^{\lambda_{h_{\nu_1}}} e_\nu)R_n(\mathbb{F};Q),$$
$$R^{\lambda}_{\beta}(\mathbb{F};Q) = R_\beta(\mathbb{F};Q)/R_\beta(\mathbb{F};Q)(\textstyle\sum_{\nu\in \mathsf{Seq}(\beta)} x_1^{\lambda_{h_{\nu_1}}} e_\nu)R_\beta(\mathbb{F};Q).$$

As a consequence of PBW theorem for the KLR algebras [Rou, Theorem 3.7], we see that $\{e_\beta \mid \mathsf{ht}(\beta)=n\}$ exhausts all the primitive central idempotents of $R_n(\mathbb{F};Q)$. It is not difficult to see that both $R^{\lambda}_{n}(\mathbb{F};Q)$ and $R^{\lambda}_{\beta}(\mathbb{F};Q)$ are finite dimensional $\mathbb{F}$-algebras.

Definition 6.8 ([Rou, §3.2.3])**.** *Let* $A=(a_{ij})_{i,j\in I}$ *be a symmetrizable GCM with the symmetrization* $d=(d_i)_{i\in I}$*. Take* $Q^A = (Q^A_{ij}(u,v))\in\mathsf{Mat}_I(\mathbb{F}[u,v])$ *subject to*

$$Q^A_{ii}(u,v)=0,\quad Q^A_{ij}(u,v)=Q^A_{ji}(v,u),\quad t_{i,j,-a_{ij},0}=t_{j,i,0,-a_{ij}}\neq 0$$

for all $i,j\in I$ *where* $Q^A_{ij}(u,v)=\sum_{\substack{p,q\geq 0\\ pd_i+qd_j=-d_ia_{ij}}} t_{ijpq}u^pv^q$.

For $n\geq 0$ and $\lambda,\beta\in\mathbb{N}[I]$ with $\mathsf{ht}(\beta)=n$, all of $R_n(\mathbb{F};Q^A)$, $R_\beta(\mathbb{F};Q^A)$, $R^{\lambda}_{n}(\mathbb{F};Q^A)$, $R^{\lambda}_{\beta}(\mathbb{F};Q^A)$ are $\mathbb{Z}$-graded via the assignment where $\nu\in I^n, 1\leq p\leq n, 1\leq a<n$.

$$\mathsf{deg}(e_\nu)=0,\quad \mathsf{deg}(x_pe_\nu)=2d_{\nu_p},\quad \mathsf{deg}(\tau_ae_\nu)=-d_{\nu_a}a_{\nu_a,\nu_{a+1}}.$$

Theorem 6.9 ([KKa])**.** *Let* $\mathbb{F}$ *be a field and let* A *be a symmetrizable GCM. Then, we can put a* $U^{\mathscr{A}}_v(A)$*-module structure on* $\bigoplus_{n\geq 0}\mathsf{K}_0(\mathsf{Proj}_{\mathsf{gr}}(R^{\lambda}_{n}(\mathbb{F};Q^A)))$ *and it is isomorphic to* $V(\lambda)^{\mathscr{A}}$.

6.3. **Graded Cartan determinants of the symmetric groups.** Recall that $\mathsf{Irr}(\mathsf{Mod}(\mathbb{Q}\mathfrak{S}_n))$ is identified with $\mathsf{Par}(n)$. For $\lambda\in\mathsf{Par}(n)$ we denote by χ_λ the corresponding character. It is characterized by $p_\mu=\sum_{\lambda\in\mathsf{Par}(n)}\chi_\lambda(C_\mu)s_\lambda$ in the ring of symmetric functions $\Lambda=\bigoplus_{n\geq 0}\varprojlim_{m\geq 0}\mathbb{Z}[x_1,\cdots,x_m]^{\mathfrak{S}_m}_n$. Recall also that any field is a splitting field for $\mathfrak{S}_n$.

Definition 6.10. *Let* $\ell\geq 2$ *be an integer.*

(a) *For each* $n\geq 0$*, we denote by* $\mathsf{Bl}_\ell(n)$ *the set of tuple* (ρ,d) *where* ρ *is an* ℓ*-core and* $d\geq 0$ *is an integer such that* $|\rho|+\ell d=n$.

(b) *For* $(\rho,d)\in\mathsf{Bl}_\ell(n)$*, we denote by* $B_{\rho,d}(\subseteq\mathsf{Par}(n))$ *the set of partition* λ *of* n *whose* ℓ*-core of* λ *is* ρ.

The following is well-known classification of p-blocks of the symmetric groups due to Nakayama-Brauer-Robinson.

Theorem 6.11. *Let* $p\geq 2$ *be a prime and let* $\lambda,\mu\in\mathsf{Par}(n)$ *be partitons of* $n\geq 0$*. The ordinary characters* χ_λ *and* χ_μ *belong to the same* p*-block if and only if* λ *and* μ *have the same* p*-core.*

Definition 6.12. *Let* $p\geq 2$ *be a prime. For* $(\rho,d)\in\mathsf{Bl}_p(n)$*, we define*

(a) $e_{\rho,d}=\sum_{\lambda\in B_{\rho,d}}\frac{\chi_\lambda(1)}{n!}\sum_{g\in\mathfrak{S}_n}\chi_\lambda(g)g^{-1}$,

(b) $\beta_{\rho,d}=\sum_{x\in\rho}\mathsf{res}(x)+d\cdot\sum_{y\in\mathbb{Z}/p\mathbb{Z}}y\in\mathbb{N}[\mathbb{Z}/p\mathbb{Z}]$ *where* $\sum_{x\in\rho}$ *means that* x *runs all the boxes of* ρ *and* $\mathsf{res}(x)=j-i+p\mathbb{Z}$ *when* x *is located at* i*-th row and* j*-th column.*

Remark 6.13. $e_{\rho,d}$ is defined as an element of $\mathbb{Q}\mathfrak{S}_n$. Thanks to Theorem 6.11, we can regard it as an element of $\mathbb{F}_p\mathfrak{S}_n$ and it is a primitive central idempotent of $\mathbb{F}_p\mathfrak{S}_n$.

Definition 6.14. *For $\ell \geq 2$, we define $Q_\ell \in \mathsf{Mat}_{\mathbb{Z}/\ell\mathbb{Z}}(\mathbb{Z}[u,v])$ by*

$$(Q_\ell)_{i,j} = \begin{cases} -(u-v)^2 & (\ell = 2 \text{ and } i \neq j) \\ \pm(v-u) & (\ell \geq 3 \text{ and } j = i \pm 1) \\ 1 & (\ell \geq 3 \text{ and } j \neq i, i \pm 1) \\ 0 & (\text{otherwise}). \end{cases} \tag{6.1}$$

For a field $\mathbb{F}$, we denote by $Q_\ell^{\mathbb{F}} \in \mathsf{Mat}_{\mathbb{Z}/\ell\mathbb{Z}}(\mathbb{F}[u,v])$ the natural image of Q_ℓ.

Theorem 6.15 ([Rou, BK2])**.** *Let $\mathbb{F}$ be a field of characteristic $p > 0$. Then, as $\mathbb{F}$-algebras we have $e_{\rho,d}\mathbb{F}\mathfrak{S}_n e_{\rho,d} \cong R^{\Lambda_0}_{\beta_{\rho,d}}(\mathbb{F}; Q_p^{\mathbb{F}})$ for $(\rho, d) \in \mathsf{Bl}_p(n)$. Especially, we have $\mathbb{F}\mathfrak{S}_n \cong R^{\Lambda_0}_n(\mathbb{F}; Q_p^{\mathbb{F}})$.*

It is known that the grading comes from Theorem 6.15 quantizes Ariki's categorification $\bigoplus_{n\geq 0} \mathsf{K}_0(\mathsf{Proj}(\mathbb{F}\mathfrak{S}_n)) \cong V(\Lambda_0)^{\mathbb{Z}}$ [BK1]. Moreover, we have a compatibility of the Shapovalov form and the graded Cartan pairing.

Theorem 6.16 ([BK3, Theorem 4.18])**.** *Let $\mathbb{F}$ be an algebraically closed field and let $\ell \geq 2$. For any $\lambda \in \mathcal{P}^+$ of $A^{(1)}_{\ell-1}$, there exist $U_v^{\mathscr{A}}(A^{(1)}_{\ell-1})$-module isomorphisms δ and ε which makes*

$$\begin{tikzcd}
V(\lambda)^{\mathscr{A}} \arrow[r, "\sim", "\delta"'] \arrow[d, "a"'] & \oplus_{n\geq 0} \mathsf{K}_0(\mathsf{Proj}_{\mathsf{gr}}(R_n^\lambda)) \arrow[d, "b"] \\
V(\lambda)^{\mathscr{A},*} & \oplus_{n\geq 0} \mathsf{K}_0(\mathsf{Mod}_{\mathsf{gr}}(R_n^\lambda)) \arrow[l, "\sim"', "\varepsilon"]
\end{tikzcd}$$

a commutative diagram. Moreover, the following diagram

$$\begin{tikzcd}
V(\lambda)^{\mathscr{A}} \otimes_{\mathscr{A}} V(\lambda)^{\mathscr{A},*} \arrow[r, "\mathsf{RSh}|_{V(\lambda)^{\mathscr{A}} \times V(\lambda)^{\mathscr{A},*}}"] \arrow[d, "\delta\otimes\varepsilon^{-1}"'] & \mathscr{A} \arrow[d, equal] \\
\oplus_{n\geq 0} \mathsf{K}_0(\mathsf{Proj}_{\mathsf{gr}}(R_n^\lambda)) \otimes_{\mathscr{A}} \oplus_{n\geq 0} \mathsf{K}_0(\mathsf{Mod}_{\mathsf{gr}}(R_n^\lambda)) \arrow[r, "\oplus\omega_{R_n^\lambda}"'] & \mathscr{A}
\end{tikzcd}$$

commutes. Here we abbreviate $R_n^\lambda(\mathbb{F}; Q_\ell^{\mathbb{F}})$ to R_n^λ and

(i) *$V(\lambda)^{\mathscr{A},*} = \{v \in V(\lambda) \mid \forall w \in V(\lambda)^{\mathscr{A}}, \langle v, w\rangle_X \in \mathscr{A}\}$ where $X \in \{\mathsf{Sh}, \mathsf{QSh}, \mathsf{RSh}\}$. The right hand side does not depend on the choice of $X \in \{\mathsf{Sh}, \mathsf{QSh}, \mathsf{RSh}\}$.*
(ii) *a is the canonical inclusion $V(\lambda)^{\mathscr{A}} \hookrightarrow V(\lambda)^{\mathscr{A},*}$.*
(iii) *b is a natural map induced from the forgetful functor $\mathsf{Proj}_{\mathsf{gr}}(R_n^\lambda) \to \mathsf{Mod}_{\mathsf{gr}}(R_n^\lambda)$.*

Remark 6.17. The presentation of Theorem 6.16 is misleading. A crucial point of the proof is that R_n^λ is isomorphic to the Ariki-Koike algebra or its degeneration [BK2, Rou].

Definition 6.18. *Let $\ell \geq 2$. We define $C^v_{\ell,d} = \mathsf{QSh}^{\mathsf{M}}_{\Lambda_0, w\Lambda_0 - d\delta}(A^{(1)}_{\ell-1})$ for $w \in W(A^{(1)}_{\ell-1})$ and $d \geq 0$. $C^v_{\ell,d}$ does not depend on the choice of w as in Proposition 4.18 and $C^v_{\ell,d}$ is related with $\mathsf{RSh}^{\mathsf{M}}_{\Lambda_0, w\Lambda_0 - d\delta}(A^{(1)}_{\ell-1})$ in the sense of Proposition 4.16 and Corollary 4.17.*

By combining Proposition 4.16, Corollary 4.17, Theorem 5.6 and Theorem 6.16, we get the graded Cartan determinants of the symmetric groups.

Theorem 6.19. *Let $\mathbb{F}$ be a filed of characteristic $p > 0$. For each $(\rho, d) \in \mathsf{Bl}_p(n)$, we have* $\det C^v_{e_{\rho,d}\mathbb{F}\mathfrak{S}_n e_{\rho,d}} = \det C^v_{\ell,d} = \prod_{s=1}^{d} [p]_s^{N_{p,d,s}}$ *where*

$$N_{p,d,s} = \sum_{\lambda \in \mathsf{Par}(d)} \frac{m_s(\lambda)}{p-1} \prod_{u \geq 1} \binom{m_u(\lambda) + p - 2}{m_u(\lambda)} = \sum_{(\lambda_i)_{i=1}^{p-1} \in \mathsf{Par}_{p-1}(d)} m_s(\lambda_1).$$

6.4. Graded Cartan determinants of the Iwahori-Hecke algebras of type A.

Definition 6.20. *Let R be a commutative domain and take $q \in R$. The Iwahori-Hecke algebra of type A with a quantum parameter q is an R-algebra $\mathcal{H}_n(R; q)$ generated by $\{T_i \mid 1 \leq i < n\}$ and obeys the following defining relations.*

$$(T_b + 1)(T_b - q) = 0, \quad T_a T_{a+1} T_a = T_{a+1} T_a T_{a+1}, \quad T_b T_c = T_c T_b$$

where $1 \leq a \leq n-2$ and $1 \leq b, c < n$ with $|b - c| > 1$.

Definition 6.21. *For $\ell \geq 2$, we fix a field k_ℓ which has an ℓ-th primitive root of unity η_ℓ.*

Remark 6.22. For $\ell \geq 2$ and $n \geq 0$, it is known that

(i) k_ℓ is a splitting field for $\mathcal{H}_n(k_\ell; \eta_\ell)$ (see [Don, §2.2]),

(ii) $\mathsf{Bl}_\ell(n)$ parametrizes the set of all primitive central idempotents of $\mathcal{H}_n(k_\ell; \eta_\ell)$.

Concerning Remark 6.22 (ii), for $(\rho, d) \in \mathsf{Bl}_\ell(n)$ we denote by $e'_{\rho,d}$ the corresponding primitive central idempotent of $\mathcal{H}_n(k_\ell; \eta_\ell)$ as in [DJ, §5]. The choice is the same as in [Rou, BK2] based on the Jucy-Murphy elements.

Theorem 6.23 ([Rou, BK2])**.** *Let $\ell \geq 2$. Then, as k_ℓ-algebras we have $e'_{\rho,d}\mathcal{H}_n(k_\ell; \eta_\ell)e'_{\rho,d} \cong R^{\Lambda_0}_{\beta_{\rho,d}}(k_\ell; Q^{k_\ell}_\ell)$ for $(\rho, d) \in \mathsf{Bl}_p(n)$. Especially, we have $\mathcal{H}_n(k_\ell; \eta_\ell) \cong R^{\Lambda_0}_n(k_\ell; Q^{k_\ell}_\ell)$.*

By the same arguments as for Theorem 6.19, we get the following.

Theorem 6.24. *For $\ell \geq 2$ and $(\rho, d) \in \mathsf{Bl}_\ell(n)$, we have* $\det C^v_{e'_{\rho,d}\mathcal{H}_n(k_\ell;\eta_\ell)e'_{\rho,d}} = \prod_{s=1}^{d} [\ell]_s^{N_{\ell,d,s}}$.

Remark 6.25. Let $\ell \geq 2$ and assume $\operatorname{char} k_\ell = 0$. In virtue of the graded version of Ariki's theorem [BK3, Corollary 5.15] and the graded Brauer-Humphreys reciprocity [HM, Theorem 2.17], we have $C^v_{e'_{\rho,d}\mathcal{H}_n(k_\ell;\eta_\ell)e'_{\rho,d}} = {}^{\mathrm{tr}}D_{\rho,d}D_{\rho,d}$ where $D_{\rho,d} = (d_{\lambda,\mu}(v))_{\lambda \in B_{\rho,d}, \mu \in B_{\rho,d} \cap \mathsf{RP}_\ell(d)}$ and $\mathsf{RP}_\ell(d)$ is the set of ℓ-regular partitions of d. Recall that we have the Fock space representation $\mathcal{F} = \bigoplus_{\lambda \in \mathsf{Par}} \mathbf{k}|\lambda\rangle$ of $U_v(A^{(1)}_{\ell-1})$ due to Hayashi and $\bigsqcup_{n \geq 0}\{G(\mu) := \sum_{\lambda \in \mathsf{Par}(n)} d_{\lambda,\mu}(v)|\lambda\rangle \mid \mu \in \mathsf{RP}_\ell(n)\}$ is Lusztig's canonical basis (also known as Kashiwara's lower global basis) of $V(\Lambda_0) \cong U_v(A^{(1)}_{\ell-1})|\phi\rangle$ (see [LLT, §6]).

7. Conjectures on the graded Cartan invarinats of the symmetric groups

7.1. Generalized blocks of the symmetric groups.

In their paper [KOR], Külshammer-Olsson-Robinson initiated the study of generalized blocks for finite groups.

Definition 7.1 ([KOR, §1])**.** *Let G be a finite group and let $\mathscr{C} \subseteq G$ be a subset of G invariant under conjugation, i.e., a union of conjugacy classes of G. Let $\mathcal{C}(G)$ be the set of complex-valued class functions on G with scalar product $\langle \alpha, \beta \rangle_{\mathscr{C}} = \frac{1}{|G|}\sum_{g \in \mathscr{C}} \alpha(g)\overline{\beta(g)}$.*

(a) *For two ordinary characters ψ, φ of G, we say that ψ and φ are directly $\mathscr{C}$-linked (and denote by $\psi \approx_{\mathscr{C}} \varphi$) if $\langle \psi, \varphi \rangle_{\mathscr{C}} \neq 0$. Clearly, the relation $\approx_{\mathscr{C}}$ is symmetric.*

(b) *A $\mathscr{C}$-block of G is an equivalence class of the ordinary characters of G under the equivalence relation $\simeq_{\mathscr{C}}$ which is defined as the transitive closure of the relation $\approx_{\mathscr{C}}$.*

Remark 7.2. For $G_{p'} := \{g \in G \mid \mathsf{ord}_G(g) \notin p\mathbb{Z}\}$ where $\mathsf{ord}_G(g)$ is the order of g in G, the notion of $G_{p'}$-blocks coincides with the usual notion of p-blocks.

Definition 7.3. *Let $\ell \geq 2$ be an integer.*

(a) *A partition λ is called ℓ-class regular if no parts of λ is divisible by ℓ (i.e., $m_{k\ell}(\lambda) = 0$ for all $k \geq 1$). We denote by $\mathsf{CRP}_\ell(n)$ the set of ℓ-class regular partitions of n.*

(b) *An ℓ-block of $\mathfrak{S}_n$ is defined as a $\mathscr{C}_{\ell'}$-block for $\mathscr{C}_{\ell'} = \bigsqcup_{\lambda \in \mathsf{CRP}_\ell(n)} C_\lambda (\subseteq \mathfrak{S}_n)$ where C_λ is a conjugacy class of $\mathfrak{S}_n$ consists of elements whose cycle type is λ.*

Theorem 7.4 ([KOR, Theorem 5.13])**.** *Let $\ell \geq 2$ be an integer and let $\lambda, \mu \in \mathsf{Par}(n)$ be partitons of $n \geq 0$. The ordinary characters χ_λ and χ_μ (see §6.3) belong to the same ℓ-block if and only if λ and μ have the same ℓ-core.*

7.2. **Conjectures of Külshammer-Olsson-Robinson and Hill.** For each $\mathscr{C}$-block B of G (see Definition 7.1) where $\mathscr{C}$ is closed (i.e., $\langle x \rangle = \langle y \rangle$ implies $y \in \mathscr{C}$ for any $x \in \mathscr{C}$ and $y \in G$), Külshammer-Olsson-Robinson assigned the Cartan group.

Definition 7.5 ([KOR, §1])**.** *Let G be a finite group and let $\mathscr{C}$ be a union of conjugacy classes of G which is closed.*

(a) *We define two additive subgroups $\mathcal{R}_G(\mathscr{C})$ and $\mathcal{P}_G(\mathscr{C})$ of $\mathcal{C}(G)$ by*

$$\mathcal{R}_G(\mathscr{C}) = \mathsf{span}_{\mathbb{Z}}\{\chi^{\mathscr{C}} \mid \chi \in \mathsf{Irr}(\mathbb{C}G)\},$$
$$\mathcal{P}_G(\mathscr{C}) = \mathcal{R}_G(\mathscr{C}) \cap \mathsf{span}_{\mathbb{Z}}\{\chi \mid \chi \in \mathsf{Irr}(\mathbb{C}G)\}$$

where $\mathsf{Irr}(\mathbb{C}G)$ is the set of ordinary characters of G and $f^{\mathscr{C}} \in \mathcal{C}(G)$ is the class function that takes the same value as f on $\mathscr{C}$ and vanishes elsewhere for $f \in \mathcal{C}(G)$.

(b) *The Cartan group $\mathsf{Cart}_G(\mathscr{C})$ is $\mathcal{R}_G(\mathscr{C})/\mathcal{P}_G(\mathscr{C})$ which is a finite group.*

We call the invariant factors of $\mathsf{Cart}_G(\mathscr{C})$ the generalized Cartan invariants. When $\ell = p$ is a prime and $\mathscr{C} = G_{p'}$, the generalized Cartan invariants are the usual Cartan invariants (i.e., the elementary divisors of the Cartan matrix $C_{\overline{\mathbb{F}_p}G}$). Recall that in the p-modular representation theory, Cartan invariants carry an information of certain centralizer groups.

Theorem 7.6 ([BrNe, Part III,§16])**.** *Let G be a finite group and let $p \geq 2$ be a prime. Take a set of representatives $g_1, \cdots, g_k$ of the p-regular conjugacy classes $C_1, \cdots C_k$ of G. The Cartan invariants of $\overline{\mathbb{F}_p}G$ is given by the multiset $\{|\,\mathsf{Syl}_p(C_G(g_i))| \mid 1 \leq i \leq k\}$.*

In the case of $G = \mathfrak{S}_n$ and $\mathscr{C} = \mathscr{C}_{\ell'}$ for $\ell \geq 2$, we may identify the Cartan group as $\mathsf{Coker}(\mathbb{Z}^{\oplus|\mathsf{RP}_\ell(n)|} \to \mathbb{Z}^{\oplus|\mathsf{RP}_\ell(n)|}, \boldsymbol{x} \mapsto \boldsymbol{x}C_{\mathcal{H}_n(k_\ell;\eta_\ell)})$ because of the following.

Theorem 7.7 ([Don, §2.2])**.** *Let $\ell \geq 2$. For any $n \geq 0$, there exist isomorphisms θ_n and ζ_n such that the following diagram commutes.*

$$\begin{tikzcd}
\mathsf{K}_0(\mathsf{Proj}(\mathcal{H}_n(k_\ell;\eta_\ell))) \arrow[r, "\sim", "\theta_n"'] \arrow[d] & \mathcal{P}_{\mathfrak{S}_n}(\mathscr{C}_{\ell'}) \arrow[d, hook] \\
\mathsf{K}_0(\mathsf{Mod}(\mathcal{H}_n(k_\ell;\eta_\ell))) \arrow[r, "\sim", "\zeta_n"'] & \mathcal{R}_{\mathfrak{S}_n}(\mathscr{C}_{\ell'})
\end{tikzcd}$$

Definition 7.8. *Let R be a commutative ring with 1.*

(a) *We say that two* $m \times m$ *matrices* $X, Y \in \mathsf{Mat}_m(R)$ *are unimodular equivalent over* R *(and denote by* $X \equiv_R Y$*) if there exist* $P, Q \in \mathrm{GL}_m(R)$ *such that* $X = PYQ$.

(b) *For a matrix* X *and a multiset* S *both consist of elements of* R*, we abbreviate* $X \equiv_R \mathsf{diag}(S)$ *to* $X \equiv_R S$ *where* $\mathsf{diag}(S) = (s_{ij})_{i,j \in I}$ *is a diagonal matrix with a multiset identity* $\{s_{ii} \mid i \in I\} = S$.

Definition 7.9. *Let* $n \geq 1$ *and let* Π *be a subset of the set of all prime numbers.*

(a) *We define* $\mathsf{pdiv}(n)$ *to be the set of prime divisors of* n *(when* $n = 1$, $\mathsf{pdiv}(n) = \emptyset$*).*

(b) *We define* n_Π *to be the* Π*-part of* n*, i.e., the unique positive integer* n_Π *such that* $n \in n_\Pi \mathbb{Z}$ *and* $\mathsf{pdiv}(n_\Pi) \subseteq \Pi, \mathsf{pdiv}(n/n_\Pi) \cap \Pi = \emptyset$.

For $\ell \geq 2$ and $(\rho, d) \in \mathsf{Bl}_\ell(n)$, we put $C_{\ell,d} := C^v_{\ell,d}|_{v=1}$ (see Definition 6.18).

Conjecture 7.10 ([KOR, Conjecture 6.4])**.** Let $\ell \geq 2$ be an integer. We define

$$r_\ell(\lambda) = \prod_{k \in \mathbb{N}_+ \setminus \ell\mathbb{Z}} (\ell/\mathsf{gcd}(\ell,k))^{\lfloor \frac{m_k(\lambda)}{\ell} \rfloor} \cdot \lfloor \frac{m_k(\lambda)}{\ell} \rfloor !_{\mathsf{pdiv}(\ell/\mathsf{gcd}(\ell,k))}$$

for a partition λ. Then, for $n \geq 0$, we have

$$\bigoplus_{(\rho,d) \in \mathsf{Bl}_\ell(n)} C_{\ell,d} \equiv_{\mathbb{Z}} \{r_\ell(\lambda) \mid \lambda \in \mathsf{CRP}_\ell(n)\}.$$

Remark 7.11. For $n \geq 0$, we may regard $\bigoplus_{(\rho,d) \in \mathsf{Bl}_\ell(n)} C^v_{\ell,d}$ as

$$\mathsf{GM}_{\mathbf{k},\tau,\mathscr{A}} \left(\oplus_{\substack{\mu \in P(\lambda) \\ \mathsf{ht}(\Lambda_0 - \mu) = n}} V(\lambda)_\mu, \oplus_{\substack{\mu \in P(\lambda) \\ \mathsf{ht}(\Lambda_0 - \mu) = n}} \mathsf{QSh}, V(\lambda)^{\mathscr{A}}_\mu \right) = \oplus_{\substack{\mu \in P(\lambda) \\ \mathsf{ht}(\Lambda_0 - \mu) = n}} \mathsf{QSh}^{\mathsf{M}}_{\Lambda_0,\mu}(A^{(1)}_{\ell-1}).$$

Remark 7.12. Let $p \geq 2$ be a prime. Then,

(a) p-regular conjugacy classes of $\mathfrak{S}_n$ is given by $\{C_\lambda \mid \lambda \in \mathsf{CRP}_p(n)\}$,

(b) the centralizer group $C_{\mathfrak{S}_n}(g)$ for $g \in C_\lambda$ is given by $\bigoplus_{k \geq 1} (\mathbb{Z}/k\mathbb{Z}) \wr \mathfrak{S}_{m_k(\lambda)}$.

Let $\ell = \prod_{p \in \mathsf{pdiv}(\ell)} p^r$ be a prime factorization of an integer $\ell \geq 2$. Hill settled affirmatively the KOR conjecture when $r \leq p$ for each $p \in \mathsf{pdiv}(\ell)$ [Hil, Theorem 1.3]. In the course of the proof, Hill proposed a following refinent of the KOR conjecture into each ℓ-blocks for $|\mathsf{pdiv}(\ell)| = 1$.

Definition 7.13. *Let* $n \geq 1$ *and let* $\ell \geq 2$*. We define* $a_\ell(n) \geq 1$ *and* $\nu_\ell(n) \geq 0$ *to be the unique integers such that* $a_\ell(n)\ell^{\nu_\ell(n)} = n$ *and* $a_\ell(n) \notin \ell\mathbb{Z}$.

Conjecture 7.14 ([Hil, Conjecture 10.5])**.** Let $p \geq 2$ be a prime and let $r \geq 1$ be an integer. For a partition λ we define a power of p by

$$\log_p I_{p,r}(\lambda) = \sum_{n \in \mathbb{N}_+ \setminus p^r\mathbb{Z}} ((r - \nu_p(n)) m_n(\lambda) + \sum_{t \geq 1} \lfloor m_n(\lambda)/p^t \rfloor).$$

Put $\ell = p^r$. Then, for each $d \geq 0$ we have

$$C_{\ell,d} \equiv_{\mathbb{Z}} \bigsqcup_{s=1}^{d} \bigsqcup_{\lambda \in \mathsf{Par}(s)} \{I_{p,r}(\lambda)\}^{u(\ell-2,d-s)}.$$

Hill proved that his conjecture implies the KOR conjecture. In fact, they are equivalent.

Theorem 7.15 ([Hil, Theorem 1.2]). *Let $\ell \geq 2$ be an integer and define $I_\ell(\lambda) = \prod_{p\in\mathsf{pdiv}(\ell)} I_{p,\nu_p(\ell)}(\lambda)$ for a partition λ. If Conjecture 7.14 is true, then for each $d \geq 0$ we have*

$$C_{\ell,d} \equiv_{\mathbb{Z}} \bigsqcup_{s=1}^{d} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{I_\ell(\lambda)\}^{u(\ell-2,d-s)}.$$

Corollary 7.16. *Conjecture 7.10 and Conjecture 7.14 are equivalent.*

Proof. Let $\ell \geq 2$ be an integer and let us assume Conjecture 7.14 is true. Then, we see that Conjecture 7.10 is true since Bessenrodt-Hill proved the multiset identity

$$\{r_\ell(\lambda) \mid \lambda \in \mathsf{CRP}_\ell(n)\} = \bigsqcup_{(\rho,d)\in\mathsf{Bl}_\ell(n)} \bigsqcup_{s=1}^{d} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{I_\ell(\lambda)\}^{u(\ell-2,d-s)} \tag{7.1}$$

for each $n \geq 0$ [BH, Theorem 5.2].

Conversely, assume Conjecture 7.10 is true. For a prime $p \geq 2$ and an integer $r \geq 1$, we put $\ell = p^r$. It is enough to prove by induction on $w \geq 0$ thatwe have $C_{\ell,w} \equiv_{\mathbb{Z}} \bigsqcup_{s=1}^{w} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{I_{p,r}(\lambda)\}^{u(\ell-2,w-s)}$. When $w = 0$, it is trivial. Because we have

$$\{r_\ell(\lambda) \mid \lambda \in \mathsf{CRP}_\ell(\ell w)\} \equiv_{\mathbb{Z}} \bigoplus_{(\rho,d)\in\mathsf{Bl}_\ell(\ell w)} C_{\ell,d} = C_{\ell,w} \oplus \bigoplus_{\substack{(\rho',d)\in\mathsf{Bl}_\ell(\ell w)\\ d<w}} C_{\ell,d},$$

inductively we must have $C_{\ell,w} \equiv_{\mathbb{Z}} \bigsqcup_{s=1}^{w} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{I_{p,r}(\lambda)\}^{u(\ell-2,w-s)}$ in virtue of (7.1). Here we use an elementary linear algebra fact that, for a PID R and finitely generated R-modules X, Y and Y', $X \oplus Y \cong X \oplus Y'$ as R-modules implies $Y \cong Y'$. $\square$

7.3. **Gradation of Hill's conjecture.** For integers $a \geq 0$ and $b \geq 1$, we denote by $a\%b$ the remainder of a by b, namely the unique integer $0 \leq c < b$ such that $a - c \in b\mathbb{Z}$.

Conjecture 7.17. Let $p \geq 2$ be a prime and let $r \geq 1$ be an integer. We define

$$I^v_{p,r}(\lambda) = \prod_{n\in\mathbb{N}_+\setminus p^r\mathbb{Z}} \prod_{k=1}^{m_n(\lambda)} [p^{r+\nu_p(k)-\nu_p(n)}]_{a_p(k)p^{\nu_p(n)}}$$

for a partition λ. Put $\ell = p^r$. Then, for each $d \geq 0$ we have

$$C^v_{\ell,d} \equiv_{\mathscr{A}} \bigsqcup_{s=1}^{d} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{I^v_{p,r}(\lambda)\}^{u(\ell-2,d-s)}.$$

Remark 7.18. We have $I^v_{p,r}(\lambda)|_{v=1} = I_{p,r}(\lambda)$ for any $\lambda \in \mathsf{Par}$. Thus, Conjecture 7.17 implies Conjecture 7.14. Note that we do not necessarily have $I^v_{p,r}(\lambda_1) \in \mathscr{A} I^v_{p,r}(\lambda_2)$ or $I^v_{p,r}(\lambda_2) \in \mathscr{A} I^v_{p,r}(\lambda_1)$ for $\lambda_1, \lambda_2 \in \mathsf{Par}(n)$ unlike at $v = 1$.

Remark 7.19. In the setting of Conjecture 7.17, Conjecture 7.17 implies the following.

$$C^v_{\ell,d} \equiv_{\mathbb{Q}[v,v^{-1}]} \bigsqcup_{s=1}^{d} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{I^v_{p,r}(\lambda)\}^{u(\ell-2,d-s)}. \tag{7.2}$$

When $r = 1$, (7.2) is the same as [ASY, Conjecture 8.2 (i)] (see Remark 6.25). Although we could not prove (7.2), it seems tractable by an elementary method since we have $C^v_{\ell,d} \equiv_{\mathbb{Q}[v,v^{-1}]} \bigsqcup_{s=1}^{d} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{\prod_{i\geq 1} [\ell]_i^{m_i(\lambda)}\}^{u(\ell-2,d-s)}$ which follows from the proof of Proposition 2.2.

As a support of Conjecture 7.17, we prove that Conjecture 7.17 gives the graded Cartan determinants of Theorem 5.6.

Theorem 7.20. *Let $p \geq 2$ be a prime and let $r \geq 1$ be an integer. Put $\ell = p^r$. Then, for each $d \geq 0$ we have the equality*

$$\prod_{s=1}^{d} [\ell]_s^{N_{\ell,d,s}} = \prod_{s=1}^{d} \prod_{\lambda \in \mathsf{Par}(s)} I^v_{p,r}(\lambda)^{u(\ell-2,d-s)}.$$

Remark 7.21. Ando-Suzuki-Yamada proved in [ASY, Theorem 4.3] that for any $\ell \geq 2$

$$\prod_{s=1}^{d} [\ell]_s^{N_{\ell,d,s}} = \prod_{s=1}^{d} \prod_{\lambda \in \mathsf{Par}(s)} Q_\ell(\lambda)^{u(\ell-2,d-s)}. \tag{7.3}$$

where $Q_\ell(\lambda) := \prod_{n \in \mathbb{N}_+ \setminus \ell\mathbb{Z}} \prod_{k=1}^{m_n(\lambda)} [\ell^{1+\nu_\ell(k)}]_{a_\ell(k)}$. Note that $I^v_{p,r}(\lambda) = Q_{p^r}(\lambda)$ when $r = 1$. Thus, Theorem 7.20 gives another product expansion of the graded Cartan determinants of the symmetric groups conjecturally comes from nice representatives of the unimodular equivalence classes of the graded Cartan matrices (but only when $|\mathsf{pdiv}(\ell)| = 1$).

Proof. In virtue of (7.3), it is enough to prove

$$\prod_{s=1}^{d} \prod_{\lambda \in \mathsf{Par}(s)} Q_{p^r}(\lambda)^{u(\ell-2,d-s)} = \prod_{s=1}^{d} \prod_{\lambda \in \mathsf{Par}(s)} I^v_{p,r}(\lambda)^{u(\ell-2,d-s)}. \tag{7.4}$$

Since $[a^b]_c = \prod_{i=1}^{b} [a]_{ca^{i-1}}$ for $a, b, c \geq 1$ and $\nu_{p^r}(n) = \lfloor \nu_p(n)/r \rfloor, a_{p^r}(n) = a_p(n) p^{\nu_p(n) \% r}$ for $n \geq 1$, it is enough to show the following multiset identity in order to prove (7.4)

$$\begin{aligned} &\bigsqcup_{s=1}^{d} \bigsqcup_{\lambda \in \mathsf{Par}(s)} \bigsqcup_{n \in \mathbb{N}_+ \setminus p^r\mathbb{Z}} \bigsqcup_{k=1}^{m_n(\lambda)} \{a_p(k)p^t \mid \nu_p(k)\%r \leq t < r + \nu_p(k)\}^{u(\ell-2,d-s)} \\ =&\bigsqcup_{s=1}^{d} \bigsqcup_{\lambda \in \mathsf{Par}(s)} \bigsqcup_{n \in \mathbb{N}_+ \setminus p^r\mathbb{Z}} \bigsqcup_{k=1}^{m_n(\lambda)} \{a_p(k)p^t \mid \nu_p(n) \leq t < r + \nu_p(k)\}^{u(\ell-2,d-s)}. \end{aligned} \tag{7.5}$$

Clearly, (7.5) follows from the following multiset identity

$$\begin{aligned} &\bigsqcup_{\lambda \in \mathsf{Par}(s)} \bigsqcup_{n \in \mathbb{N}_+ \setminus p^r\mathbb{Z}} \bigsqcup_{k=1}^{m_n(\lambda)} \{a_p(k)p^t \mid \nu_p(n) \leq t < r + \nu_p(k)\} \\ =& \bigsqcup_{\lambda \in \mathsf{Par}(s)} \bigsqcup_{n \in \mathbb{N}_+ \setminus p^r\mathbb{Z}} \bigsqcup_{k=1}^{m_n(\lambda)} \{a_p(k)p^t \mid \nu_p(k)\%r \leq t < r + \nu_p(k)\} \end{aligned}$$

for $s \geq 1$ which follows from Proposition 7.22. ☐

Proposition 7.22. *Let $p \geq 2$ be a prime and let $r \geq 1$ be an integer. We have the following multiset identity for each $d \geq 1$ and $u \in \mathbb{N}_+ \setminus p\mathbb{Z}$.*

$$\bigsqcup_{\lambda \in \mathsf{Par}(d)} \bigsqcup_{n \geq 1} \bigsqcup_{\substack{1 \leq k \leq m_n(\lambda) \\ a_p(k) = u}} \{\nu_p(n)\} = \bigsqcup_{\lambda \in \mathsf{Par}(d)} \bigsqcup_{n \geq 1} \bigsqcup_{\substack{1 \leq k \leq m_n(\lambda) \\ a_p(k) = u}} \{\nu_p(k)\%r\}.$$

Before giving a proof, we need preparatory definitions.

Definition 7.23. *For $d \geq 1$, we define maps $\mathsf{cut}_d, \mathsf{Infl}_d : \mathsf{Par} \to \mathsf{Par}$ by the following.*

(a) *for $k \geq 1$, $m_k(\mathsf{cut}_d(\lambda)) = m_k(\lambda)$ if $k \notin d\mathbb{Z}$, otherwise $m_k(\mathsf{cut}_d(\lambda)) = 0$.*

(b) *$\ell(\lambda) = \ell(\mathsf{Infl}_d(\lambda))$ and $(\mathsf{Infl}_d(\lambda))_i = d\lambda_i$ for $1 \leq i \leq \ell(\lambda)$,*

Definition 7.24. *Let $p \geq 2$ be a prime and let $r \geq 1$ be an integer. Note that any integer $a \geq 1$ can be uniquely expressed as $a = \sum_{i=0}^{r-1} a_i p^i$ with $a_{r-1} \geq 0$ and $0 \leq a_j < p$ for $0 \leq j \leq r-2$. Define the subset $\Psi^{p,r}_a$ of $\mathsf{Par}(a)$ consists of all the partitions ν such that*

(i) *for all $0 \leq k < r$, we have $\sum_{h=k}^{r-1} a_h p^{h-k} \geq \sum_{h=k}^{r-1} m_{p^h}(\nu) p^{h-k}$,*

(ii) *we have $\sum_{h=0}^{r-1} a_h p^h = \sum_{h=0}^{r-1} m_{p^h}(\nu) p^h$.*

Remark 7.25. $\Psi^{p,r}_a$ is characterized as the smallest subset of $\mathsf{Par}(a)$ such that

(a) Define $\lambda(a) \in \mathsf{Par}(a)$ by $m_{p^i}(\lambda(a)) = a_i$ for $0 \leq i < r$. Then, we have $\lambda(a) \in \Psi^{p,r}_a$.

(b) For any $\lambda \in \Psi^{p,r}_a$ and any $0 \leq i < r$ with $m_i(\lambda) > 0$, define $\mu \in \mathsf{Par}(a)$ to be

$$m_j(\mu) = \begin{cases} m_i(\lambda) - 1 & (j = i) \\ m_{i-1}(\lambda) + p & (j = i-1) \\ m_j(\lambda) & (\text{otherwise}) \end{cases}$$

for $0 \leq j < r$. Then, we have $\mu \in \Psi^{p,r}_a$.

Thus, we have $\Psi^{p,r}_a = \{\lambda \in \mathsf{CRP}_{p^r}(a) \mid \forall f \in \mathbb{N}_+ \setminus p^{\mathbb{N}}, m_f(\lambda) = 0\}$.

Proof. Because of Lemma 7.26, it is enough to prove the following multiset identity for each $a \geq 1$ and $u \in \mathbb{N}_+ \setminus p\mathbb{Z}$.

$$\bigsqcup_{\lambda \in \Psi^{p,r}_a} \bigsqcup_{n \geq 1} \bigsqcup_{\substack{1 \leq k \leq m_n(\lambda) \\ a_p(k) = u}} \{\nu_p(n)\} = \bigsqcup_{\lambda \in \Psi^{p,r}_a} \bigsqcup_{n \geq 1} \bigsqcup_{\substack{1 \leq k \leq m_n(\lambda) \\ a_p(k) = u}} \{\nu_p(k)\% r\}. \tag{7.6}$$

For any $0 \leq s < r$, we define sets $U_1(s)$ and $U_2(s)$ as follows.

$$U_1(s) = \{(\lambda, j) \in \Psi^{p,r}_a \times \mathbb{N} \mid up^j \leq m_{p^s}(\lambda)\},$$
$$U_2(s) = \{(\lambda, j, t) \in \Psi^{p,r}_a \times \mathbb{N} \times \{0, \cdots, r-1\} \mid up^j \leq m_{p^t}(\lambda), j - s \in r\mathbb{Z}\}.$$

Note that (7.6) is the same as $\bigsqcup_{0 \leq s < r} \{s\}^{|U_1(s)|} = \bigsqcup_{0 \leq s < r} \{s\}^{|U_2(s)|}$. Thus, it is enough to construct a bijection between $U_1(s)$ and $U_2(s)$ for any $0 \leq s < r$.

We claim that the map $U_2(s) \to U_1(s), (\lambda, j, t) \mapsto (\mu, i)$ given by

$$i = t + j - s, \quad m_{p^w}(\mu) = \begin{cases} m_{p^t}(\lambda) - up^j & (s \neq t \text{ and } w = t) \\ m_{p^s}(\lambda) + up^i & (s \neq t \text{ and } w = s) \\ m_{p^w}(\lambda) & (\text{otherwise}) \end{cases}$$

where $0 \leq w < r$ is well-defined. To see this, only non-trivial check should be to prove $\mu \in \Psi^{p,r}_a$ when $s > t$. In order to prove this, it is enough to check (i) in Definition 7.24 for $\nu = \mu$ and $h = s$. Since $\lambda \in \Psi^{p,r}_a$, we have $\sum_{h=t}^{r-1} a_h p^{h-t} \geq \sum_{h=t}^{r-1} m_{p^h}(\lambda) p^{h-t}$. Thus,

$$\begin{aligned} \sum_{h=s}^{r-1} a_h p^{h-s} + p^{t-s} \sum_{h=t}^{s-1} a_h p^{h-t} &\geq \sum_{h=s}^{r-1} m_{p^h}(\lambda) p^{h-s} + p^{t-s} \sum_{h=t}^{s-1} m_{p^h}(\lambda) p^{h-t} \\ &\geq \sum_{h=s}^{r-1} m_{p^h}(\lambda) p^{h-s} + m_{p^t}(\lambda) p^{h-t} \geq \sum_{h=s}^{r-1} m_{p^h}(\mu) p^{h-s}. \end{aligned}$$

Since $0 \leq p^{t-s}\sum_{h=t}^{s-1} a_h p^{h-t} \leq p^{t-s}\sum_{h=t}^{s-1}(p-1)\cdot p^{h-t} = 1-p^{t-s} < 1$, we are done.

Similarly, we see that the map $U_1(s) \to U_2(s), (\mu, i) \mapsto (\lambda, j, t)$ given by

$$t = i\%r, \quad j = i+s-t, \quad m_{p^w}(\lambda) = \begin{cases} m_{p^t}(\mu) + up^j & (s \neq t \text{ and } w = t) \\ m_{p^s}(\mu) - up^i & (s \neq t \text{ and } w = s) \\ m_{p^w}(\mu) & (\text{otherwise}) \end{cases}$$

is well-defined. Clearly, it is a converse of the map $U_2(s) \to U_1(s)$ above. □

Lemma 7.26. *Let $p \geq 2$ be a prime and let $r \geq 1$ be an integer. For each $d \geq 1$, the multiset $S_d^{p,r} := \{\mathsf{cut}_{p^r}(\lambda) \mid \lambda \in \mathsf{Par}(d)$ such that $\mathsf{cut}_{p^r}(\lambda) \neq \phi\}$ has a decomposition of the form $S_d^{p,r} = \bigsqcup_{i\in I} \mathsf{Sort}(\prod_{j\in J_i} \mathsf{Infl}_{d_{i,j}}(\Psi^{p,r}_{a_{i,j}}))$ such that*

(a) *I is a finite non-empty set and $\{J_i \mid i \in I\}$ is a family of finite non-empty sets,*
(b) *$a_{i,j} \geq 1$ and $d_{i,j} \in \mathbb{N}_+ \setminus p\mathbb{Z}$ for $i \in I$ and $j \in J_i$,*
(c) *for any $i \in I$ and $j \neq j' \in J_i$, take $\lambda \in \mathsf{Infl}_{d_{i,j}}(\Psi^{p,r}_{a_{i,j}})$ and $\mu \in \mathsf{Infl}_{d_{i,j'}}(\Psi^{p,r}_{a_{i,j'}})$. Then, we have either $m_f(\lambda) = 0$ or $m_f(\mu) = 0$ for all $f \geq 1$,*
(d) *$\mathsf{Sort} : \mathsf{Par}^* \to \mathsf{Par}$ is a map that maps a finite sequence of partitions $(\lambda^{(g)})_{g\in G}$ to a partition μ such that $\{\mu_f \mid 1 \leq f \leq \ell(\mu)\} = \bigsqcup_{g\in G}\{\lambda^{(g)}_f \mid 1 \leq f \leq \ell(\lambda^{(g)})\}$ as multisets.*

Proof. Let λ be a non-empty partition. Clearly, $L = \{1, \cdots, \ell(\lambda)\}$ has a decomposition of the form $L = \bigsqcup_{k\in K} L_k$ such that

(i) for any $k \in K$, we have $L_k \neq \emptyset$ and there exists a unique $d_k \in \mathbb{N}_+ \setminus p\mathbb{Z}$ such that $\lambda_j \in d_k\mathbb{Z}$ and $\lambda_j/d_k \in p^{\mathbb{N}}$ for all $j \in L_k$,
(ii) for any $k \neq k' \in K$, we have $d_k \neq d_{k'}$.

Put $\lambda^{(k)} = \mathsf{Sort}(((\lambda_j/d_k))_{j\in L_k})$ for $k \in K$. As in Remark 7.25, there exists a unique $a_k \geq 1$ such that $\lambda^{(k)} \in \Psi^{p,r}_{a_k}$. Thus, we have $\lambda \in \mathsf{Sort}(\prod_{k\in K} \mathsf{Infl}_{d_k}(\Psi^{p,r}_{a_k}))$.

From this construction, we see that $\mathsf{CRP}_{p^r}(d)$ has a decomposition of the form $\mathsf{CRP}_{p^r}(d) = \bigsqcup_{i\in I} \mathsf{Sort}(\prod_{j\in J_i} \mathsf{Infl}_{d_{i,j}}(\Psi^{p,r}_{a_{i,j}}))$ that satisfies (a), (b) and (c) for each $d \geq 1$.

Because we have $S_d^{p,r} = \bigsqcup_{0\leq e<d/p^r} \mathsf{CRP}_{p^r}(d-p^r e)^{|\mathsf{Par}(e)|}$, we are done. □

7.4. Gradation of Külshammer-Olsson-Robinson's conjecture when $\ell = p^r$.

By the same reason to consider a gradation of Hill's conjecture (see §1.3), we want to consider a correct gradation of the KOR conjecture. Though we could not find it in general, we propose a gradation of the KOR conjecture only when $|\,\mathsf{pdiv}(\ell)| = 1$.

Conjecture 7.27. Let $p \geq 2$ be a prime and let $r \geq 1$ be an integer. Put $\ell = p^r$. Then, for each $n \geq 0$, we have $\bigoplus_{(\rho,d)\in\mathsf{Bl}_\ell(n)} C^v_{\ell,d} \equiv_{\mathscr{A}} \{r^v_{p,r}(\lambda) \mid \lambda \in \mathsf{CRP}_\ell(n)\}$ where

$$r^v_{p,r}(\lambda) = \prod_{k\geq 1}\prod_{t=1}^{\lfloor\frac{m_k(\lambda)}{\ell}\rfloor} [p^{r-\nu_p(k)}]_{tp^{\nu_p(k)}} \cdot [p^{\nu_p(t)}]_{a_p(t)p^{\nu_p(k)}} = \prod_{k\geq 1}\prod_{t=1}^{\lfloor\frac{m_k(\lambda)}{\ell}\rfloor} [p^{r-\nu_p(k)+\nu_p(t)}]_{a_p(t)p^{\nu_p(k)}}.$$

Remark 7.28. When $r = 1$, Conjecture 7.27 implies [ASY, Conjecture 8.2 (ii)] which predicts $\bigoplus_{(\rho,d)\in\mathsf{Bl}_p(n)} C^v_{p,d} \equiv_{\mathbb{Q}[v,v^{-1}]} \{r^v_{p,1}(\lambda) \mid \lambda \in \mathsf{CRP}_p(n)\}$ for $n \geq 0$ and a prime $p \geq 2$.

Proposition 7.29. *Conjecture 7.17 implies Conjecture 7.27. More precisely, we have the following multiset identity for each $r, n \geq 1$ and a prime $p \geq 2$ where $\ell = p^r$.*

$$\bigsqcup_{(\rho,d)\in\mathsf{Bl}_\ell(n)}\ \bigsqcup_{s=1}^{d}\ \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{I^v_{p,r}(\lambda)\}^{u(\ell-2,d-s)} = \{r^v_{p,r}(\lambda) \mid \lambda \in \mathsf{CRP}_\ell(n)\}.$$

Proof. Since $r^v_{p,r}(\lambda) = I^v_{p,r}(\mathsf{red}_\ell(\lambda))$ and $I^v_{p,r}(\lambda) = I^v_{p,r}(\mathsf{cut}_\ell(\lambda))$ for any partition λ where $m_k(\mathsf{red}_\ell(\lambda)) = \lfloor m_k(\lambda)/\ell \rfloor$ for all $k \geq 1$, Proposition 7.29 follows from Lemma 7.30. $\square$

Lemma 7.30 ([BH, Lemma 5.5])**.** *Let $\ell \geq 2$ be an integer. We have the multiset identity*

$$\bigsqcup_{(\rho,d)\in\mathsf{Bl}_\ell(n)} \bigsqcup_{s=1}^{d} \bigsqcup_{\lambda\in\mathsf{Par}(s)} \{\mathsf{cut}_\ell(\lambda)\}^{u(\ell-2,d-s)} = \{\mathsf{red}_\ell(\lambda) \mid \lambda \in \mathsf{CRP}_\ell(n)\}$$

for any $n \geq 0$.

References

[Aka] T. Akasaka, *An integral PBW basis of the quantum affine algebra of type $A^{(2)}_2$*, Publ.Res.Inst.Math.Sci. **38** (2002), 803–894.

[ASY] M. Ando, T. Suzuki and H-F. Yamada, *Combinatorics for graded Cartan matrices of the Iwahori-Hecke algebra of type A*, to appear in Ann.Comb., **arXiv:1005.1134v5**.

[APW] H. Andersen, P. Polo, K. Wen, *Representations of quantum algebras*, Invent.Math. **104** (1991), 1–59.

[Bec] J. Beck, *Braid group action and quantum affine algebras*, Comm.Math.Phys. **165** (1994), 555–568.

[BCP] J. Beck, V. Chari and A. Pressley, *An algebraic characterization of the affine canonical basis*, Duke Math.J. **99** (1999), 455–487.

[BH] C. Bessenrodt and D. Hill, *Cartan invariants of symmetric groups and Iwahori-Hecke algebras*, J.Lond.Math.Soc. **81** (2010), 113–128.

[BK1] J. Brundan and A. Kleshchev, *The degenerate analogue of Ariki's categorification theorem*, Math.Z. **266** (2010), 877–919.

[BK2] J. Brundan and A. Kleshchev, *Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras*, Invent.Math. **178** (2009), 451–484.

[BK3] J. Brundan and A. Kleshchev, *Graded decomposition numbers for cyclotomic Hecke algebras*, Adv.Math. **222** (2009), 1883–1942.

[BK4] J. Brundan and A. Kleshchev, *Cartan determinants and Shapovalov forms*, Math.Ann. **324** (2002), 431–449.

[BK5] J. Brundan and A. Kleshchev, *Hecke-Clifford superalgebras, crystals of type $A^{(2)}_{2\ell}$ and modular branching rules for $\widehat{S}_n$*, Represent.Theory **5** (2001), 317–403.

[BKM] G. Benkart, S-J. Kang and D. Melville, *Quantized enveloping algebras for Borcherds superalgebras*, Trans.Amer.Math.Soc. **350** (1998), 3297–3319.

[BeNa] J. Beck and H. Nakajima, *Crystal bases and two-sided cells of quantum affine algebras*, Duke Math.J. **123** (2004), 335–402.

[BO1] C. Bessenrodt and J. Olsson, *A note on Cartan matrices for symmetric groups*, Arch.Math. **81** (2003), 497–504.

[BO2] C. Bessenrodt and J. Olsson, *On Cartan matrices and lower defect groups for covering groups of symmetric groups*, Trans.Amer.Math.Soc. **358** (2006), 3623–3635.

[Bra] R. Brauer, *On the Cartan invariants of groups of finite order*, Ann.of Math.(2) **42** (1941), 53–61.

[BrNe] R. Brauer and C. Nesbitt, *On the modular characters of groups*, Ann.of Math.(2) **42** (1941), 556–590.

[CJ] V. Chari and N. Jing, *Realization of level one representations of $U_q(\widehat{\mathfrak{g}})$ at a root of unity*, Duke Math.J. **108** (2001), 183–197.

[DcKK] C. De Concini, V. Kac and D. Kazhdan, *Boson-fermion correspondence over $\mathbb{Z}$*, Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988), 124–137, Adv.Ser.Math.Phys., **7**, World Sci.Publ., Teaneck, NJ, 1989.

[DJ] R. Dipper and G. James, *Blocks and idempotents of Hecke algebras of general linear groups*, Proc.London Math.Soc. **54** (1987), 57–82.

[Don] S. Donkin, *Representations of Hecke algebras and characters of symmetric groups*, Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000), 49–67, Progr.Math., **210**, Birkhaüser Boston, Boston, MA, 2003.

[Dri] V. Drinfeld, *A new realization of Yangians and of quantum affine algebras*, Soviet Math.Dokl. **36** (1988), 212–216.

[FJ] I. Frenkel and N. Jing, *Vertex representations of quantum affine algebras*, Proc.Nat.Acad.Sci.U.S.A. **85** (1988), 9373–9377.

[Her] D. Hernandez, *Kirillov-Reshetikhin conjecture: the general case*, Int.Math.Res.Not. 2010, 149–193.

[Hil] D. Hill, *Elementary divisors of the Shapovalov form on the basic representation of Kac-Moody algebras*, J.Algebra **319** (2008), 5208–5246.

[HM] J. Hu and A. Mathas, *Graded cellular bases for the cyclotomic Khovanov-Lauda-Rouquier algebras of type A*, Adv.Math. **225** (2010), 598–642.

[HW] D. Hill and W. Wang, *Categorification of quantum Kac-Moody superalgebras*, **arXiv:1202.2769**.

[Jin] N. Jing, *Twisted vertex representations of quantum affine algebras*, Invent.Math. **102** (1990), 663–690.

[JM] N. Jing and K. Misra, *Vertex operators for twisted quantum affine algebras*, Trans.Amer.Math.Soc. **351** (1999), 1663–1690.

[Kac] V. Kac. *Infinite dimensional Lie algebras*. Cambridge University Press, 1990.

[Kas] M. Kashiwara, *On crystal bases*, Representations of groups (Banff, AB, 1994), 155–197, CMS Conf.Proc., **16**, Amer.Math.Soc., Providence, RI, 1995.

[KKa] S-J.Kang and M.Kashiwara *Categorification of Highest Weight Modules via Khovanov-Lauda-Rouquier Algebras*, to appear in Invent.Math.

[KKw] S-J. Kang and J-H. Kwon, *Fock space representations of quantum affine algebras and generalized Lascoux-Leclerc-Thibon algorithm*, J.Korean Math.Soc. **45** (2008), 1135–1202.

[KKT] S-J. Kang, M. Kashiwara and S. Tsuchioka, *Quiver Hecke superalgebras*, arXiv:1107.1039

[KL1] M. Khovanov and A. Lauda, *A diagrammatic approach to categorification of quantum groups. I.*, Represent.Theory **13** (2009), 309–347.

[KL2] M. Khovanov and A. Lauda, *A diagrammatic approach to categorification of quantum groups II.*, Trans.Amer.Math.Soc. **363** (2011), 2685–2700.

[KOR] B. Külshammer, J. Olsson and G. Robinson, *Generalized blocks for symmetric groups*, Invent.Math. **151** (2003), 513–552.

[Lak] V. Lakshmibai, *Bases for quantum Demazure modules*, Representations of groups (Banff, AB, 1994), 199–216, CMS Conf.Proc., **16**, Amer.Math.Soc., Providence, RI, 1995.

[LLT] A. Lascoux, B. Leclerc and J-Y. Thibon, *Hecke algebras at roots of unity and crystal bases of quantum affine algebras*, Comm.Math.Phys. **181** (1996), 205–263.

[LT] B. Leclerc and J-Y. Thibon, *q-deformed Fock spaces and modular representations of spin symmetric groups*, J.Phys.A **30** (1997), 6163–6176.

[Lus] G. Lusztig, *Introduction to quantum groups*, Reprint of the 1994 edition. Modern Birkhaüser Classics. Birkhaüser/Springer, New York, 2010.

[Lus2] G. Lusztig, *Finite dimensional Hopf algebras arising from quantized universal enveloping algebras*, J.Amer.Math.Soc. **3** (1990), 257–296.

[Ols] J. Olsson, *Lower defect groups in symmetric groups*, J. Algebra **104** (1986), 37–56.

[Rou] R. Rouquier, *2-Kac-Moody algebras*, arXiv:0812.5023

[Tsu] S. Tsuchioka, *Hecke-Clifford superalgebras and crystals of type $D_\ell^{(2)}$*, Publ.Res.Inst.Math.Sci. **46** (2010), 423–471.

Kavli Institute for the Physics and Mathematics of the Universe, Todai Institutes for Advanced Study, the University of Tokyo, Kashiwa, Japan 277-8583 (Kavli IPMU)

E-mail address: `tshun@kurims.kyoto-u.ac.jp`